A Review of Methicillin-Resistant *Staphylococcus aureus* (MRSA) on Milk and Milk Products: Public Health Importance

Aswin Rafif Khairullah¹, Sri Agus Sudjarwo², Mustofa Helmi Effendi^{3*}, Nenny Harijani⁴, Wiwiek Tyasningsih⁵, Jola Rahmahani⁵, Dian Ayu Permatasari⁴, Sancaka Cashyer Ramandinianto⁶, Agus Widodo¹, Katty Hendriana Priscilia Riwu¹

¹Doctoral Program in Veterinary Science, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia. ²Department of Veterinary Pharmacology, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia. ³Halal Research Center, Universitas Airlangga, Surabaya, Indonesia.

⁴Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia.

⁵Department of Veterinary Microbiology, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia.

⁶Master Program in Veterinary Disease and Public Health, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia.

*Corresponding Author E-mail: mheffendi@yahoo.com

ABSTRACT

Methicillin-resistant Staphylococcus aureus (MRSA) is a pathogenic strain of bacteria in humans and animals that can cause anything from mild to severe infections. Detection of MRSA in foodstuffs of animal origin has been widely reported and has raised public concerns about the transmission of MRSA from foodstuffs of animal origin to humans. There have been many reports of MRSA infections originating from dairy cow's milk, milk products, and MRSA transmission between farmers working in dairy pens and people working in the dairy industry. Significant differences in the spread of cases of MRSA infection originating from milk and milk products have been reported among various countries in the world. This difference may be caused by different livestock production management systems in different countries in the world. Milk and milk products contaminated with MRSA will be a potential transmission of MRSA transmission when consumed by humans and can be a problem for public health. The milking process carried out by farmers and the processing of milk for milk products can be risk factors for transmission of MRSA to humans. Contamination of milk and milk products by MRSA is a major cause of food poisoning. Several antibiotics of choice can be used in treating patients who experience poisoning with milk and milk products such as fluoroquinolone. tetracyclines, lincosamide, and trimethoprimsulfamethoxazole. Adherence to good hygiene practices during milking, processing and handling of dairy cow milk can significantly minimize MRSA contamination of milk and dairy products.

INTRODUCTION

Methicillin-resistant Staphylococcus aureus (MRSA) is a pathogenic strain of bacteria in humans and animals that can cause from mild to severe infections and can even cause death cases worldwide. The epidemiology of the incidence of this infection is also very diverse, considering the possible exchange of MRSA strains between humans and animals or vice versa [1]. Recently this year, the detection of MRSA in foodstuffs of animal origin has been widely reported and has raised public concerns about the transmission of MRSA from foodstuffs of animal origin to humans [2, 3]. Various types of foodstuffs of animal origin have been studied to determine the source of MRSA transmission, such as beef, chicken, pork, dairy cow's milk, sheep's milk, and dairy products [4-10]. It is widely known that Staphylococcus aureus, especially methicillinresistant Staphylococcus aureus (MRSA) is the main cause of mastitis cases in dairy cows, even MRSA can also be detected in healthy cows [11-14].

During the milking process in dairy cows, especially in dairy cows with subclinical mastitis, this MRSA strain can emerge from the teats of dairy cows to contaminate milk without organoleptic changes in milk, thus allowing this MRSA strain to easily spread to the milk processing Keywords: MRSA, milk, milk products, public health

Correspondence:

Mustofa Helmi Effendi

³Halal Research Center, Universitas Airlangga, Surabaya, Indonesia. ***Corresponding Author E-mail**: mheffendi@yahoo.com

process. There have been many reports of MRSA infections originating from dairy milk [11, 13, 15], dairy products [10], and transmission of MRSA between farmers working in dairy pens and people working in the dairy industry [16, 17].

MRSA transmission can be zoonotic, involving direct animal-to-human contact or vice versa, the processing of dairy products, and contact with environmental contaminants [18]. MRSA transmission disease cases have become a serious problem for dairy farming and the dairy industry [19-21].

Poor management of dairy farms and inappropriate use of antibiotics can increase the incidence of MRSA in cow's milk [22]. Previous studies have reported cases of hospital-related MRSA infection (Hospital acquired Methicillin-resistant Staphylococcus aureus / HA-MRSA) Community acquired Methicillin-resistant and Staphylococcus aureus (CA-MRSA) infections. Recent studies have found another group of MRSA, namely MRSA with livestock (Livestock associated associated Methicillin-resistant Staphylococcus aureus / LA-MRSA). LA-MRSA colonization of dairy cows has been shown to be a risk factor for veterinarians, breeders, employees of the

dairy processing industry, and those in close contact with dairy cows [23].

LA-MRSA infections in humans, originating from milk and milk products, can include skin and soft tissue infections ranging from mild to severe [24]. Expression of methicillin-resistant Staphylococcus aureus may complicate antibiotic treatment [25]. The prevalence of LA-MRSA needs to be done from the beginning before the transmission of food from animal origin to humans can eventually become a threat to human health [26]. Studies on the spread of MRSA that are sourced from foodstuffs of animal origin such as milk and dairy products can limit the threat of MRSA infection cases to human health [23].

This review describes Methicillin-resistant Staphylococcus aureus in general, MRSA in foodstuffs of animal origin, MRSA in milk, MRSA in milk products, epidemiology of MRSA in milk and milk products, transmission of MRSA transmission from milk and dairy products, risk factors. transmission of MRSA originating from milk and milk products, consequences for public health, treatment of cases of MRSA poisoning originating from milk and milk products, and prevention of MRSA poisoning from milk and milk products.

Methicillin-resistant Staphylococcus aureus (MRSA)

Expression of resistance to penicillinase-stable antibiotics to penicillinase, commonly referred to as "methicillin resistance" or "oxacillin resistance", in staphylococcus aureus bacteria is manifested as bacteria that are resistant to all β -lactam class antibiotic agents including carbapenems and sephalosporins [27].

The start of methicillin-resistant Staphylococcus aureus (MRSA) began in 1961 in the United Kingdom [28], one year after methicillin antibiotics were introduced into clinical practice to treat cases of penicillin-resistant Staphylococcus aureus infection [29]. In the early 90's MRSA began to become a serious health threat in the United Kingdom and the United States [30]. Resistance is encoded by a mobile genetic element, known as staphylococcal chromosomal cassette mec (SCCmec) [31], carrying the genes encoding mecA and mecC, both of which encode penicillin-binding protein 2a (PBP2a) [28, 32]. B-lactam antibiotics bind to PBP, acting in the synthesis of cell wall peptidoglycan, which causes bacterial cells to lysis. But PBP2a has a low affinity for βlactam antibiotics, so that peptidoglycan synthesis activity continues in MRSA strains [32], despite the presence of various β-lactam inhibitor combinations. The mecA coding gene can be identified in one of eleven SCCmec (I-XI) types, which carry 5 different types of mec-coding gene complexes, such as mecA and their regulatory genes mecR1 and mecI, and eight different types of the ccr gene complex., which contains two different recombinase CCR genes that play a role in the mobility of the element [33]. MRSA was obtained from the integration of the SCCmec element carrying the mecA gene encoding the methicillinsensitive Staphylococcus aureus [34]. A new MRSA coding gene named mecC (originally named mecALGA251) by the International Working Group (IWG) was discovered in the SCCmec element classification, which was isolated for the first time from a sample of dairy cow milk in the UK in 2011 [35] and this mecC coding gene was also isolated from human clinical samples in Denmark, Scotland, and the UK in 2014 [36]. This mecC gene is located in the SCCmec type XI element and has 69% similarity in the identity of the nucleotide sequence to the mecA coding gene. Besides MRSA strains have the ability to adapt to the

pressure of β -lactam antibiotics, MRSA strains also have the potential to produce enterotoxins that can cause foodborne disease. However, the prevalence of enterotoxigenic MRSA in food and foodstuffs is still low and is usually closely related to milk and milk products. In a previous study conducted in Italy, it was reported that three MRSA strains (1.2%) isolated from milk and dairy products were found to have enterotoxigenic properties [37], and in the United States study 2 MRSA strains (1.33%) were found. those isolated from tank milk had enterotoxins [12].

Methicillin-resistant *Staphylococcus aureus* (MRSA) in food of animal origin

Many researchers who have reported finding MRSA in foodstuffs of animal origin around the world, have found varying prevalences in beef, lamb, chicken, pork, rabbit, wild boar, eggs, milk, dairy products, and fish [1, 4, 5, 6, 8, 10, 38-43]. In addition, it is important to emphasize that the MRSA strains found in humans have also been found in food, due to contamination of food consumed by humans [44]. It also depends on the epidemiological factors of the geographic area of each country, about 0.7-1.5% of humans in the world have been infected with MRSA [45-47].

Until now it is known that there is a transmission of MRSA clone exchange between humans and animals [48], but it is not clear whether there is a risk of inherited foodstuffs of animal origin associated with MRSA contamination. In 1994 there was a case of food poisoning from food of animal origin at the University of Rotterdam hospital, where out of 27 patients 5 of them died, the incident was caused by an isolated strain of MRSA from a cook contaminating food during food preparation [49]. Also, in 2001 there was a case of food poisoning from food of animal origin caused by MRSA strains contained in Staphylococcus enterotoxin C, which was isolated from roasted pork contaminated by food handlers [50].

MRSA in milk

The incidence of MRSA infection from cow isolates with mastitis has been widely studied and the prevalence rate is still low [51]. Following the first reports of MRSA from cows with mastitis [52], sporadic cases of MRSA infection in dairy cows have been detected among Staphylococcus aureus isolates from dairy cows with subclinical mastitis and clinical mastitis. In a study conducted in Korea [53], MRSA of cow's milk was found with an isolation ratio of 0.18%. In a report on dairy farming in Belgium, the highest percentage of MRSA identification was found in breastfeeding dairy cows at 15% [51], these cows did have a previous history of MRSA infection. Until now, the prevalence of MRSA identification in dairy cows with mastitis is still low, considering the time period since MRSA was first discovered in dairy cows and close contact with farmers with dairy cows' udders. In a study in Germany, the highest proportion of positive MRSA samples was found in the nose swabs of calves, namely 45% and the lowest proportion of positive samples of MRSA in bulk tank milk was 4.1%, most of the MRSA isolates came from the spa type t011 and t034 belonging to the CC398 clonal complex [4]. The discovery of LA-MRSA CC398 in tank milk proved that MRSA was colonized in dairy cows' udders and a possible cause of subclinical mastitis cases in dairy cows in Germany [4]. Close contact between dairy cows and humans can lead to strain transfer between dairy cows and humans through milk production. In a study in Hungary, identical MRSA isolates were found

from dairy cows with mastitis and a farmer with phenotypic and genotype analysis which indicated that there had been a transfer of MRSA strains between dairy cows and humans [54].

MRSA in milk products

Staphylococcus aureus and MRSA can contaminate milk either by direct excretion from udders of dairy cows with mastitis or through contamination from the surrounding environment during handling and processing of milk for milk products. It is known that milk is a suitable medium for bacterial development and a means of transmission of bacterial transmission such as MRSA when milk or milk products are consumed by humans, so this is a public health hazard [55].

One of the livestock-based food products that can be a source of MRSA transmission and cause cases of food poisoning by MRSA is milk from dairy cows, then milk from dairy cows that is contaminated with MRSA is processed into milk products. MRSA strains have been identified from a wide variety of dairy products such as cheese [8], yogurt [56], butter [57], ice cream [58], and pasteurized milk [59]. This MRSA strain also possibly carries the Staphylococcal enterotoxin gene which could potentially cause food poisoning [8].

Epidemiology MRSA in milk and milk products

The cause of transmission of MRSA to humans is obtained from milk cows and milk products contaminated with MRSA [8, 12]. Consumption or handling of milk and dairy products can lead to the spread of the antibiotic resistance gene Staphylococcus aureus to humans [60].

The spread of MRSA infection cases originating from milk and milk products has been reported in various countries around the world. Different livestock production management systems in different countries of the world. Different policies and regulations for the use of antibiotics in each country also contribute to the different spread rates of MRSA infection [61]. Several animal species in an area that cause the transfer of genetic material between MRSA strains can affect the rate of spread of MRSA infection [17]. Reported data explain the wide variety of spread of MRSA infections in milk and dairy products in various parts of the world.

Europe

In a French study, no MRSA was found among 119 Staphylococcus aureus isolates in the milk of dairy cows with clinical mastitis between 1998 and 2000 [62]. In a similar study in France, only 1 Staphylococcus aureus isolate was classified as MRSA among 139 Staphylococcus aureus isolates in dairy cattle with clinical mastitis between 2007 and 2008 [63]. In a large study conducted in Hungary from January 2002 to December 2004, 139 Staphylococcus aureus isolates (out of 595 samples of cow's milk with subclinical mastitis) detected 27 of them MRSA [54].

Based on a survey conducted in Belgium between 2006 and 2007, out of 118 Staphylococcus aureus isolates obtained from milk samples from dairy cows with clinical mastitis, 11 were confirmed to be MRSA [13].

Research conducted in the Czech Republic between 2006 and 2009, [64] found 18 MRSA strains from cow's bulk tank milk and 4 MRSA strains from goat's milk from a total of 299 Staphylococcus aureus isolates studied. One of the MRSA isolates from the goat farm was detected positive for the gene encoding Staphylococcal enterotoxin (SE) [64]. Moreover, between 2006 and 2008, Stastkova et al. [65] found 5 MRSA strains from 34 isolates of Staphylococcus aureus obtained from 153 milk samples from goat farms, and all these MRSA isolates contained the Staphylococcal enterotoxin B (seb) gene. The lower prevalence of MRSA reported in the Czech Republic is 3%, in this study there were 60 samples of goat's milk, 60 samples of sheep's milk, and 120 samples of dairy cow's milk, only 2 were detected strains of MRSA, 2 of which were dairy cows with mastitis. subclinical from one herd [66].

A study conducted in Italy [8] found 4 MRSA strains carrying staphylococcal enterotoxins from cow's milk samples and dairy products (mozzarella cheese and pecorino cheese), while other researchers [67] did not find MRSA strains from 36 Staphylococcus aureus isolates extracted from cheese sample made from sheep's milk.

A study conducted in Portugal [68] did not find any MRSA among 30 Staphylococcus aureus isolates from milk samples from dairy cows with subclinical mastitis, whereas a study conducted in Ireland [60] did not find MRSA strains with mecA positive in 70 milk samples that were not pasteurized. A study conducted in Slovakia [69] did not find MRSA strains from 79 isolates of Staphylococcus aureus obtained from sheep's milk samples and cheese samples made from sheep's milk.

In Switzerland [70] found only 2 MRSA strains among 142 Staphylococcus aureus isolates obtained from milk samples from dairy cows with mastitis. A survey conducted in Germany on tank milk samples between 2009 and 2010 described an estimated prevalence of MRSA infection cases of 1% to 2% [71] and 5% [72], respectively. According to a study conducted in southwest Germany on three dairy farms, a substantial prevalence of MRSA was found to be 5% - 17% of dairy samples and 1% - 10% of dairy cattle [17]. In a study conducted in the Netherlands [73] detected 62 MRSA strains from 1,839 dairy samples collected from 26 dairy farms.

America

In the United States, [74] reported a MRSA prevalence of 1% of 846 Staphylococcus aureus isolates obtained from dairy samples on Michigan farms, another study [75] reported a MRSA prevalence of 2% of 2132 Staphylococcus aureus isolates obtained from dairy samples collected from 1994 to 2001 on a Wisconsin farm. No MRSA strains were found among the 357 Staphylococcus aureus isolates obtained from milk samples taken from 24 dairy herds on farms in North Carolina and Virginia [76]. In a survey research conducted by Virgin et al. [77], no MRSA strains were found in 542 bulk tank milk samples. D'amico and Donnelly [78] reported that have isolated 2 MRSA strains among 90 Staphylococcus aureus isolates obtained from raw milk to be used for the production of handmade cheese in Vermont. Haran et al. [12] isolated 2 MRSA strains from 154 Staphylococcus aureus isolates obtained from 150 bulk tank milk samples on a Minnesota farm. They also reported that the two MRSA strains were capable of producing enterotoxins such as SEB, SEC, SED and SEE. In Brazil, there was a high MRSA prevalence rate of 25% of the 98 milk samples of dairy cows with subclinical mastitis [79], however in a Canadian study, Saini et al. [80] found only one MRSA strain among 1,802 Staphylococcus aureus isolates obtained from milk samples collected from 79 dairy farms.

Asia

In a study in Japan between 1998 and 2005 [81] found 4 MRSA strains from 363 isolates of Staphylococcus aureus obtained from dairy cows from 260 dairy farms.

In a study in Pakistan [82] revealed an MRSA prevalence of 10%, there were 8 positive MRSA strains out of 77 Staphylococcus aureus isolates obtained from dairy cow's milk.

In a study in Korea between May 2001 and April 2003 [83] revealed an MRSA prevalence of 1.5%, there were 12 MRSA positive strains from 894 samples of dairy milk. Meanwhile, the survey conducted [53] between 1999 and 2003, there was a prevalence of less than 0.5% from examining 9,055 samples of milk from dairy cows. Kwon et al. [53] revealed that all identified MRSA strains carried the Staphylococcal enterotoxin gene (SED, SEI, SEJ). On the other hand, according to two surveys conducted between 2003 and 2009 [84] and between 1997 and 2004 [85], there are estimates of MRSA prevalence of 2.5% and 6% on examinations of 835 and 402 samples of dairy cows with mastitis.

In a study conducted in Turkey, there was an MRSA prevalence of 17.5%, 18 positive MRSA strains were found among 103 Staphylococcus aureus isolates from milk samples from dairy cows with mastitis [86]. In the latest survey results [87] did not find MRSA strains among 21 Staphylococcus aureus isolates obtained from 857 sheep milk samples and 33 goat milk samples collected from 13 farms in Kirikkale province.

In a study conducted in India between 2007 and 2008 [88] revealed a MRSA prevalence rate of 10% among 128 Staphylococcus aureus isolates obtained from milk samples from Karan Fries (Taurus × Zebu) cows suffering from mastitis. Meanwhile, the same research team [89] has also found a MRSA prevalence rate of 13% among 107 Staphylococcus aureus isolates from a herd of dairy cows on farms in northwest India.

In a survey research conducted in Thailand in 2010 [90] identified 74 MRSA strains out of 375 isolates of Staphylococcus aureus obtained from 598 milk samples from dairy cows with mastitis, 376 bulk tank milk samples, and 46 pasteurized milk samples.

In a study conducted in Iran in 2010 [58] examined 300 samples (100 raw milk, 100 ice cream, 100 pasteurized milk) and found 20 MRSA strains carrying the mecA-coding gene (14 from the raw milk sample and 6 from the ice cream sample). Whereas in a study conducted between 2010 and 2011, examined 348 samples of randomly selected dairy cows and reported an MRSA prevalence rate of 28% [91].

Africa

In a study conducted in South Africa [92] reported an MRSA prevalence rate of 6% of dairy cow's milk produced on 2 commercial farms, and in a study conducted in Ethiopia [93] found a high MRSA prevalence rate of 60% of 78 Staphylococcus aureus isolates obtained from dairy cows.

In a study conducted in Nigeria, Suleiman et al. [94] examined 339 milk samples from dairy cows and reported an MRSA prevalence rate of 36%, found 26 isolates of Staphylococcus aureus resistant to oxacillin and 2 of them carrying the mecA gene coding.

Transmission of MRSA

MRSA strains can spread from animals to humans and from humans to animals. MRSA transmission usually

occurs through direct contact, often through contact with animals or people who have been infected or colonized with MRSA [83, 95].

The level of spread of MRSA in the general population usually varies depending on the geographical conditions of each country [96]. In the hospital, patients who have been infected and colonized with MRSA will be the main reservoir of transmission of MRSA, which is usually easily spread from patient to patient by hand touch [97]. The route of transmission of MRSA may be similar to that of other strains of Staphylococcus aureus, but there may be differences in the efficiency of host colonization after exposure [98]. There are many other factors in the transmission of MRSA transmission such as through contaminated surrounding air, saliva splashes, and nasal discharge [99-101].

Animals also act as reservoirs for MRSA transmission and can transmit MRSA to humans and other animals [102-107]. Several types of MRSA exhibit host specificity and are widely spread over certain geographic areas. Therefore, MRSA transmission is more frequently associated with animals than with humans [108-110]. LA-MRSA CC398 is the type of MRSA that is most associated with intensive reared animal-based food-producing livestock, especially dairy cows [97]. Although LA-MRSA CC398 was found to colonize livestock without causing clinical symptoms, it could cause clinical infection with MRSA in animals. LA-MRSA CC398 colonization has been found in many breeders, family members of farmers, employees of the food processing industry, and veterinarians [96, 111, 112]. The spread of MRSA can also occur in personnel and other animals including cats, dogs, horses, between pets, in veterinary hospitals, and in households [44, 113-116]. In some cases, pets can act as a source of MRSA infection in humans [95, 117].

Foodstuffs of animal origin including milk and milk products contaminated with MRSA and the handling process that contaminates MRSA in milk and milk products are also potential transmission transmissions. In the case of patients hospitalized due to food poisoning, MRSA contamination can spread MRSA to other patients or healthcare workers [97]. Reports in the Netherlands, MRSA cases related to contaminated milk infected 27 patients and 14 hospital workers and caused 5 deaths. A person who consumed milk and milk products was found to have colonized the MRSA strains similar to the MRSA strains found in milk and milk product samples. Contaminated milk and milk products may lead to cases of septicemia due to MRSA, which can then be passed on to other patients in the hospital. Airborne transmission that is contaminated with MRSA also plays an important role in the spread of MRSA [49].

Given the abundance of evidence for the presence of MRSA strains in dairy cows, it will raise public concern about MRSA contaminants found in milk and dairy products [118]. However, there is no detailed description of the direct association of MRSA in animal and human foodstuffs, although there have been reports of increased MRSA contamination in milk and milk products, as well as the incidence of MRSA infection in the community [97]. Further investigations are needed to determine the true role of milk and dairy products in animal-to-human transmission of MRSA transmission.

Another aspect of MRSA that contaminates milk and dairy products is that MRSA often carries Staphylococcal enterotoxin (SE) genes such as SEA, SEB, SEC, SED which are often the cause of food poisoning (EFSA, 2008). Different combinations of MRSA clones are associated with the staphylococcal enterotoxin gene, but the reasons for this association have not been widely elucidated [119, 120]. Clinically, cases of food poisoning caused by MRSA are almost the same as cases of food poisoning caused by other strains of Staphylococcus aureus [118]. MRSA strains that contaminate milk and dairy products can also be of human origin, due to less aseptic handling and processing systems for milk and dairy products [50]. The increasing prevalence of MRSA strains among the Staphylococcus aureus strains may lead to an increase in the prevalence of Staphylococcus aureus toxicity [121].

Risk Factor of MRSA

Studies conducted in the United States, in the human population about 30% of humans are infected with methicillin-susceptible Staphylococcus aureus (MSSA) [122] and about 1.5% of humans are infected with methicillin-resistant Staphylococcus aureus [45, 123] with MSSA colonization sites. and MRSA mostly occurs in the nostrils [124]. While such colonization does not cause clinical symptoms in an infected person, it may be a risk factor for developing subsequent symptomatic infection [122, 125]. Humans infected with MRSA without showing symptoms are categorized as persistent carriages and intermittent carriages that act as MRSA carriers and can transmit MRSA to more susceptible persons [124].

Staphylococcus aureus is a common bacterial agent causing mastitis in dairy cows. A study conducted in Minnesota to estimate the prevalence rate of Staphylococcus aureus in bulk tank milk reported that the MSSA and MRSA prevalence rates were 84% and 4%, respectively [12, 125]. The results of another study estimated that the prevalence rate of Staphylococcus aureus in bulk milk tanks was 35% in samples of dairy cows in Louisiana [12] and 31% in samples of dairy cows in Pennsylvania. Studies from Ireland [126], Turkey [127], Argentina [128], and Brazil [129] have found the presence of the Staphylococcal enterotoxin gene in MRSA derived from cow's milk. Udder of dairy cows with clinical and subclinical mastitis may play a role in the contamination of milk by MRSA via direct excretion of the cow's nipples [130] with large fluctuations in the count of zero to 108 CFU / ml [131]. For example, mastitis cases in dairy cows were the only source of contamination in 1999 in Brazil, then MRSA from cow's milk affected 328 people who consumed unpasteurized dairy cow's milk [132]. Likewise, 293 Staphylococcus aureus isolates were isolated from 127 goat milk samples and sheep milk samples in Switzerland [130]. Recently, MRSA strains were found in the milk of dairy cows with mastitis and bulk tank milk on Hungarian farms which suggest that udders of dairy cows infected with MRSA can contaminate milk and milk products [131]. However, MRSA contamination of milk and milk products can occur in the environment during the processing and handling of milk [124]. Therefore, the milking process carried out by the farmer, the processing and handling of milk for milk products can be a risk factor for the transmission of MRSA to humans [132].

Public Health Importance

Contamination of milk and milk products by MRSA is a major cause of food poisoning. May be characterized by gastroenteritis symptoms which are clinically manifested as emesis with or without accompanied diarrhea. These cases of poisoning stem from ingestion of one or more Staphylococcal enterotoxins that are formed in milk and

milk products that have been contaminated with MRSA. Symptoms of systemic toxicity, such as hypotension and fever are rare cases of MRSA-related food poisoning. Symptoms of systemic toxicity resolve on their own within 24-48 hours after onset. The exact incidence is still not widely known, but MRSA is probably the most common cause of food poisoning in the United States [133]. The risk assessment in milk and milk products relies on detection of classic bacteria and counting the number of coagulase-positive Staphylococcus on selective Baird-Parker media. It is necessary to observe the progression of the disease after consumption of milk and dairy products, the presence of toxic cells, and the production of toxic cells. Therefore, in some countries, low levels of contamination by Staphylococcus aureus and MRSA are tolerated in most food products (e.g. 103 CFU / g of Staphylococcus aureus contamination in French cheese) is not usually considered a public health risk. [131, 133].

Treatment of food-borne intoxication MRSA from milk and milk products

In cases of poisoning due to consuming milk and milk products contaminated with MRSA, several antibiotics can be used in treating patients who have poisoned milk and milk products. Penicillin is usually still used as the antibiotic of choice as long as Staphylococcus aureus infection is sensitive to it [134]. Semisynthetic penicillins, such as methicillin are indicated for patients infected with β-lactamase-producing Staphylococcus aureus [135, 136]. MRSA-infected patients are treated with a glycopeptide known as the antibiotic vancomycin [136, 137]. Vancomycin is the main antibiotic of choice in the treatment of cases of MRSA infection [136, 137]. Patients who are allergic to vancomycin antibiotics can be given alternative antibiotics such as fluoroquinolone (ciprofloxacin), tetracyclines (minocycline), lincosamide (clindamycin), trimethoprim-sulfamethoxazole also known as cotrimoxazole [135, 136]. Recent quinolones, such as the antibiotic ciprofloxacin which has increased anti-Staphylococcal activity can be used as antibiotics of choice, but their use is still limited because of the rapid development of Staphylococcus aureus resistance during treatment [135, 136]. MRSA infection cases can also be treated with several oral drugs such as rifampicin-fusidic acid, linezolid, rifampicin fluoroquinolone, + cotrimoxazole (trimethoprim-sulphamethoxazole), doxycycline or minocyline, pristinamycin, and clindamycin [138]. Mupirocin (Bactroban) also has the potential to fight MRSA from mucous membrane colonization [139]. A recent antibiotic called platensimycin has also shown success in curing cases of MRSA infection [140].

Prevention of food-borne intoxication MRSA from milk and milk products

The increasing number of cases of poisoning of milk and milk products due to MRSA strains requires proper precautions. These precautions must be taken to prevent contamination throughout the food processing, production chain and preparation [141]. Adhering to good hygiene practices during milking, processing, and handling of dairy milk can significantly minimize contamination of milk and milk products. Increased vigilance in maintaining cleanliness must be carried out by everyone in the preparation and processing of dairy cow milk so that the safety quality of milk and milk products can be maintained [141, 142]. Milk must also be properly heated to make it safe for consumption. The results of a recent analysis of growth requirements revealed that Staphylococcus aureus and MRSA can grow at a water activity of 0.867 and at temperatures as low as 8 ° C [143]. Therefore, it is necessary to provide counseling for industrial workers, breeders and the general public regarding the dangers of MRSA contamination in milk and milk products. There are also many cases of milk poisoning by Staphylococcus aureus and MRSA because the milk is left at room temperature for too long, for example during picnics or large banquets, so that the agenda gives Staphylococcus aureus and MRSA to develop and produce Staphylococcal enterotoxin (SE) 143].

The human health risks associated with consumption of unpasteurized milk and milk products [144, 145, 146]. However, Staphylococcal enterotoxin produced from MRSA is very resistant to heating and pasteurization, heating milk for one hour can reduce the amount of toxins, but only autoclave at 15 psi for 20 minutes can completely destroy the toxin. Milk that has been sterilized still requires cooling at a temperature of 0 ° C to 4 ° C so that later the milk can be processed further. Since MRSA is known to grow well on salt media, there is a higher risk of MRSA contamination in home salted cheese [147].

Conclusion

Milk and milk products are potential vehicles for MRSA transmission when consumed by humans. The association between the multidrug-resistance nature of MRSA and the enterotoxigenic properties presents a serious public health risk, because enterotoxins can be found in milk and milk products contaminated with MRSA. Hygiene monitoring in the processing and handling of milk needs to be done continuously in order to obtain the quality of milk and milk products that are hygienic and safe for consumption. For this reason, the application of good manufacturing practices and hazard analysis critical control point (HACCP) is very important to ensure the quality and safety of milk and milk products.

Acknowledgements

This review was supported in part with the **Penelitian Hibah Mandat** funding from Airlangga University, Indonesia in the fiscal year 2020, with grant number: 368/UN3.14/PT/2020.

REFERENCES

- 1. Parisi A, Caruso M, Normanno G, Latorre L, Sottili R, Miccolupo A, et al. Prevalence, antimicrobial susceptibility and molecular typing of Methicillinresistant *Staphylococcus aureus* (MRSA) in bulk tank milk from southern Italy. Food Microbiol. 2016; 58: 36-42.
- 2. EFSA (European Food Safety Authority). Scientific opinion of the panel on biological hazards on a request from the European commission on Assessment of the public health significance of methicillin-resistant *Staphylococcus aureus* (MRSA) in animal and foods. EFSA J. 2009; 993: 1-73.
- 3. Larsen J, Stegger M, Andersen PS, Petersen A, Larsen AR, Westh H, et al. Evidence for human adaptation and foodborne transmission of livestock-associated methicillin-resistant *Staphylococcus aureus*. Clin Infect Dis. 2016; 63: 1349-52.

- 4. Tenhagen BA, Vossenkuhl B, Kasbohrer A, Alt K, Kraushaar B, Guerra B, et al. Methicillin-resistant *Staphylococcus aureus* in cattle food chains prevalence, diversity, and resistance in Germany. J Anim Sci. 2014; 92: 2741-51.
- 5. Fessler A, Kadlec K, Hassel M, Hauschild T, Eidam C, Ehricht R, et al. Characterization of methicillinresistant *Staphylococcus aureus* isolates from food and food products of poultry origin in Germany. Appl Environ Microbiol. 2011; 77(20): 7151-7.
- 6. de Boer E, Zwartkruis-Nahuis JTM, Wit B, Huijsdens XW, de Neeling AJ, Bosch T, et al. Prevalence of methicillin-resistant *Staphylococcus aureus* in meat. Int J Food Microbiol. 2009; 134: 52-6.
- 7. O'Brien AM, Hanson BM, Farina SA, Wu JY, Simmering JE, Wardyn SE, et al. MRSA in conventional and alternative retail pork products. PloS One. 2012; 7: e30092.
- Normanno G, Corrente M, La Salandra G, Dambrosio A, Quaglia NC, Parisi A, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) in foods of animal origin product in Italy. Int J Food Microbiol. 2007; 117: 219-22.
- 9. Carfora V, Giacinti G, Sagrafoli D, Marri N, Giangolini G, Alba P, et al. Methicillin-resistant *Staphylococcus aureus* and methicillin-susceptible *Staphylococcus aureus* in dairy sheep and incontact humans: an intra-farm study. J Dairy Sci. 2006; 99: 4251-8.
- Basanisi MG, La Bella G, Nobili G, Franconieri I, La Salandra G. Genotyping of methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from milk and dairy products in South Italy. Food Microbiol. 2017; 62: 141-6.
- Fessler A, Scott C, Kadlec K, Ehricht R, Monecke S, Schwarz S. Characterization of methicillin-resistant *Staphylococcus aureus* ST398 from cases of bovine mastitis. J Antimicrob Chemother. 2010; 65: 619-25.
- Haran KP, Godden SM, Boxrud D, Jawahir S, Bender JB, Sreevastan S. Prevalence and characterization of *Staphylococcus aureus*, including methicillin-resistant *Staphylococcus aureus*, isolated from bulk tank milk from Minnesota dairy farms. J Clin Microbiol. 2012; 50: 688-95.
- 13. Vanderhaeghen W, Cerpentier T, Adriaensen C, Vicca J, Hermans K, Butaye P. Methicillin-resistant *Staphylococcus aureus* (MRSA) ST398 associated with clinical and subclinical mastitis in Belgian cows. Vet Microbiol. 2010; 144: 166-71.
- 14. Pu W, Su Y, Li J, Li C, Yang Z, Deng HP, et al. High incidence of oxacillin-susceptible *mec*A-positive *Staphylococcus aureus* (OS-MRSA) associated with bovine mastitis in China. PloS One. 2014; 9(2): e88134.
- 15. Benedetti V, Cremonesi P, Ferrari S, Castiglioni B, Fabbi M, Vicari N, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) from bovine milk samples. Large Anim Rev. 2010; 16: 67-70.
- Antoci E, Pinzone MR, Nunnari G, Stefani S, Cacopardo B. Prevalence and molecular characteristics of methicillin-resistant *Staphylococcus aureus* (MRSA) among subjects working on bovine dairy farms. Infez Med. 2013; 21: 125-9.
- Spohr M, Rau J, Friedrich A, Klittich G, Fetsch A, Guerra B, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) in three dairy herds in southwest Germany. Zoonoses Public Health. 2011; 58: 252-61.

- 18. Nunang C, Young R. MRSA in farm animals and meat-A new threat to human health. Soil association, USA. 2007.
- 19. Elhaig MM, Selim A, Mahmoud MM, El-Gayar EK. Molecular confirmation of *Trypanosoma evansi* and *Babesia bigemina* in cattle from lower Egypt. Pak Vet J. 2016; 36: 409-14.
- Qayyum A, Khan JA, Hussain R, Awais M, Ahmad N, Khan MS. Investigation of milk and blood serum biochemical profile as an indicator of subclinical mastitis in Cholistani cattle. Pak Vet J. 2016; 36: 275-9.
- 21. Yilmaz R, Cangul IT, Onat K, Akkoc A, Ozyigit MO, Akdesir E. Histopathological, immunohistochemical and bacteriological characterization of *Mycoplasma bovis* pneumonia in cattle. Pak Vet J. 2016; 36: 316-21.
- 22. Joshi LR, Tiwari A, Devkota SP, Khatiwada S, Paudyal S, Pande KR. Prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) in dairy farms of Pokhara, Nepal. Int J Vet Sci. 2014; 3: 87-90.
- Paterson GK, Larsen J, Harrison EM, Larsen AR, Morgan FJ, Peacock SJ, et al. First detection of livestock-associated meticillin-resistant *Staphylococcus aureus* CC398 in bulk tank milk in the United Kingdom, January to July 2012. Euro surveillance: bulletin Europeen sur les maladies transmissibles, European communicable disease bulletin. 2012; 17(50).
- 24. Layer F, Cuny C, Strommenger B, Werner G, Witte W. AktuelleDaten und Trends zu. Methicillin-resistenten *Staphylococcus* aureus (MRSA). Bundesgesundheitsbla. 2012; 55: 1377-86.
- 25. Lowy FD. Antimicrobial resistance: the example of *Staphylococcus aureus*. J Clin Invest. 2003; 111: 1265-73.
- Kadlec K, Ehricht R, Monecke S, Steinacker U, Kaspar H, Mankertz J, et al. Diversity of antimicrobial resistance pheno-and genotypes of methicillinresistant *Staphylococcus aureus* ST398 from diseased swine. J Antimicrob Chemother. 2009; 350.
- 27. Stapleton PD, Taylor PW. Methicillin resistance in *staphylococcus aureus*. Sci Prog. 2002; 85(1): 57-72.
- 28. Petinaki E, Spiliopoulou I. Methicillin-Resistant *Staphylococcus aureus* among Companion and Food-Chain Animals: Impact of Human Contact. Clin Microbiol Infect. 2012; 18: 626-34.
- 29. Jevons MP. Celbenin-Resistant Staphylococci. Br Med J. 1961; 1(5129): 124-5.
- 30. Kluytmans J. Methicillin-Resistant *Staphylococcus aureus* in Food Products: cause for concern or case for complacency? Clin Microbial Infect. 2010; 16: 11-5.
- Milheiriço C, Oliveira DC, de Lencastre H. Multiplex PCR Strategy for Subtyping the Staphylococcal Cassette 492 Chromosome *mec* type IV in Methicillin-Resistant *Staphylococcus aureus*: 'SCCmec IV multiplex. J Antimicrob Chemoth. 2007; 60: 42-8.
- 32. Paterson GK, Morgan FJE, Harrison EM, Peacok SJ, Parkhill J, Zadoks RN, et al. Prevalence and Properties of *mecC* Methicillin-Resistant *Staphylococcus aureus* (MRSA) in Bovine Tank Milk in Great Britain. J Antimicrob Chemoth. 2014; 69: 598-602.
- 33. Kondo Y, Teruyo I, Ma XX, Watanabe S, Kreiswirth BN, Etienne J, et al. Combination of Multiplex PCRs for Staphylococcal Cassette Chromosome *mec* Type Assignment: Rapid Identification System for *mec, ccr*, and Major Differences in Junkyard Regions. Antimicrob Agents Chemother. 2007; 51: 264-74.

- 34. Enright MC, Robinson DA, Randle G, Feil EJ, Grundmann H, Spratt BG. The Evolutionary History of Methicillin Resistant *Staphylococcus aureus* (MRSA). Proc Natl Acad Sci. 2002; 99: 7687-92.
- 35. García-Álvarez L, Holden MTG, Lindsay H, Webb CR, Brown DFJ, Curran MD, et al. Methicillin-Resistant *Staphylococcus aureus* with a Novel *mecA* Homologue in Human and Bovine Populations in the UK and Denmark: A descriptive study. Lancet Infect Dis. 2011; 11: 595-603.
- van Duijkeren E, Hengeveld PD, Albers M, Pluister G, Jacobs P, Heres L, et al. Prevalence of Methicillin-Resistant *Staphylococcus aureus* Carrying *mecA* or *mecC* in Dairy Cattle. Vet Microbiol, 2014; 171: 364-7.
- 37. Carfora V, Caprioli A, Marri N, Sagrafoli D, Boselli C, Giacinti G, et al. Enterotoxin Genes, Enterotoxin Production, and Methicillin Resistance in *Staphylococcus aureus* Isolated from Milk and Dairy Products in Central Italy. Int Dairy J. 2015; 42: 12-5.
- Doulgeraki AI, Di Ciccio P, Lanieri A, Nychas GJE. Methicillin-resistant food-related *Staphylococcus aureus*: a review of current knowledge and biofilm formation for future studies and applications. Res Microbiol. 2017; 168(1): 1-15.
- Parisi A, Caruso M, Normanno G, Latorre L, Miccolupo A, Fraccalvieri R, et al. High occurrence of methicillinresistant *Staphylococcus aureus* in horses at slaughterhouses compared with those for recreational activities: a professional and food safety concern? Foodborne Pathog Dis. 2017; 14(12): 735-41.
- 40. Normanno G, Dambrosio A, Lorusso V, Samoilis G, Di Taranto P, Parisi A. Methicillin-resistant *Staphylococcus aureus* (MRSA) in slaughtered pigs and abattoir workers in Italy. Food Microbiol. 2015; 51: 51-6.
- 41. Syed MA, Shah SHH, Sherafzal Y, Shafi-Ur-Rehman S, Khan MA, Barrett JB, et al. Detection and molecular characterization of methicillin-resistant *Staphylococcus aureus* from table eggs in Haripur, Pakistan. Foodborne Pathog Dis. 2018; 15(2): 86-93.
- 42. Mohammad MO, Salman AEB, Lafi SQ. Prevalence of *Staphylococcus aureus* in imported fish and correlations between antibiotic resistance and enterotoxigenicity. J Food Protect. 2015; 78(11): 1999-2005.
- 43. Wendlandt S, Schwarz S, Silley P. Methicillin-Resistant *Staphylococcus aureus*: A Food-Borne Pathogen? Annu Rev Food Sci T. 2013; 4: 117-39.
- 44. Weese JS, Dick H, Willey BM, McGeer A, Kreiswirth BN, Innis B, et al. Suspected transmission of methicillinresistant *Staphylococcus aureus* between domestic pets and humans in veterinary clinics and in the household. Vet Microbiol. 2006; 115: 148-55.
- 45. Gorwitz RJ, Kruszon-Moran D, McAllister SK, McQuillan G, McDougal LK, Fosheim GE, et al. Changes in the prevalence of nasal colonization with *Staphylococcus aureus* in United States, 2001-2004. J Infect Dis. 2008; 197(9): 1226-34.
- 46. Wertheim HFL, Vos MC, Boelens HAM, Voss A, Vandenbroucke-Grauls CMJE, Meester MHM, et al. Low prevalence of methicillin resistant *Staphylococcus aureus* (MRSA) at hospital admission in The Netherlands: the value of search and destroy and restrictive antibiotic use. J Hosp Infect. 2004; 56(4): 321-25.

- 47. Munckhof WJ, Nimmo GR, Schooneveldt JM, Schlebusch S, Stephens AJ, Williams G, et al. Nasal carriage of *Staphylococcus aureus*, including community-associated methicillin-resistant strains, in Queensland adults. Clin Microbiol Infect. 2009; 15(2): 149-55.
- 48. Zarazaga M, Gomez P, Ceballos S, Torres C. Molecular epidemiology of *Staphylococcus aureus* lineages in the animal-human interface. Editor: Alexandra Fetsch, *Staphylococcus aureus*, Academic Press. 2018; 189-214.
- 49. Kluytmans J, van Leeuwen W, Goessens W, Hollis R, Messer S, Herwaldt L, et al. Food initiated outbreak of methicillin-resistant *Staphylococcus aureus* analyzed by phenotyping and genotyping. J Clin Microbiol. 1995; 33(5): 1121-8.
- 50. Jones TF, Kellum ME, Porter SS, Bell M, Schaffner W. An outbreak of community-acquired foodborne illness caused by methicillin-resistant *Staphylococcus aureus*. Emerg Infect Dis. 2002; 8(1): 82-4.
- 51. Hendriksen RS, Mevius DJ, Schroeter A, Teale C, Meunier D, Butaye P, et al. Prevalence of antimicrobial resistance among bacterial pathogens isolated from cattle in different European countries: 2002-2004. Acta Vet Scand. 2008; 50: 28.
- 52. Devriese LA, Van Damme LR, Fameree L. Methicillin (cloxacillin)-resistant *Staphylococcus aureus* strains isolated from bovine mastitis cases. Zentralbl Veterinarmed B. 1972; 19(7): 598-605.
- 53. Kwon NH, Park KT, Moon JS, Jung WK, Kim SH, Kim JM, et al. Staphylococcal cassette chromosome *mec* (SCC*mec*) characterization and molecular analysis for methicillin-resistant *Staphylococcus aureus* and novel SCC*mec* subtype IVg isolated from bovine milk in Korea. J Antimicrob Chemother. 2005; 56: 624-32.
- 54. Juhasz-Kaszanyitzky E, Janosi S, Somogyi P, Dan A, van der Graafvan Bloois L, van Duijkeren E, et al. MRSA transmission between cows and humans. Emerg Infect Dis. 2007; 13: 630-2.
- 55. Le Jeune JT, Schultz PJR. Unpasteurized milk: a continued public health threat. Clin Infect Dis. 2009; 48: 93-100.
- 56. Papadopoulos P, Papadopoulos T, Angelidis AS, Boukouvala E, Zdragas A, Papa A, et al. Prevalence of *Staphylococcus aureus* and of methicillin-resistant *Staphylococcus aureus* (MRSA) along the production chain of dairy products in north-western Greece. Food Microbiol. 2018; 69: 43-50.
- 57. Titouche Y, Hakem A, Houali K, Hennekinne JA, Torres C, Auvray F. Emergence of methicillin-resistant *Staphylococcus aureus* (MRSA) ST8 in raw milk and traditional dairy products in the Tizi Ouzou area of Algeria. J Dairy Sci. 2019; 102(8): 6876-84.
- 58. Mirzaei H, Farhoudi H, Tavassoli H, Farajli M, Monadi A. Presence and antimicrobial susceptibility of methicillin resistant *Staphylococcus aureus* in raw and pasteurized milk and ice cream in Tabriz by culture and PCR techniques. Afr J Microbiol Res. 2012; 6: 6224-9.
- 59. Dai J, Wu S, Huang J, Wu Q, Zhang F, Zhang J, et al. Prevalence and characterization of *Staphylococcus aureus* isolated from pasteurized milk in China. Front Microbiol. 2019; 10: 641.
- McKay AM. Antimicrobial resistance and heat sensitivity of oxacillin-resistant, *mecA*-positive *Staphylococcus spp*. from unpasteurized milk. J Food Protect. 2008; 71: 186-90.

- Grave K, Torren-Edo J, Mackay D. Comparison of the sales of veterinary antibacterial agents between 10 European countries. J Antimicrob Chemother. 2010; 65: 2037-40.
- 62. Guerin-Faublee V, Carret G, Houffschmitt P. In vitro activity of 10 antimicrobial agents against bacteria isolated from cows with clinical mastitis. Vet Rec. 2003; 152: 466-71.
- 63. Haenni M, Galofaro L, Ponsin C, Bes M, Laurent F, Madec JY. Staphylococcal bovine mastitis in France: enterotoxins, resistance and the human Geraldine methicillin-resistant *Staphylococcus aureus* clone. J Antimicrob Chemother. 2011; 66: 216-25.
- 64. Stastkova Z, Karpiskova S, Karpiskova R. Findings of methicillin-resistant strains of *Staphylococcus aureus* in livestock. Czech J Food Sci. 2009; 27: 236-41.
- 65. Stastkova Z, Karpiskova S, Karpiskova R. Occurrence of methicillin-resistant strains of *Staphylococcus aureus* at a goat breeding farm. Veterinarni Medicina. 2009; 54: 419-26.
- 66. Vyletělova M, Hanuš O, Karpíšková R, Štástková, Z. Occurrence and antimicrobial sensitivity in staphylococci isolated from goat, sheep and cow's milk. Acta Universitatis Agric Silvic Mendelianae Brunensis. 2011; 59: 209-13.
- 67. Spanu V, Virdis S, Scarano C, Cossu F, De Santis EPL, Cosseddu AM. Antibiotic resistance assessment in *Staphylococcus aureus* strains isolated from raw sheep's milk cheese. Vet Res Commun. 2010; 34: 87-90.
- 68. Nunes SF, Bexiga R, Cavaco LM, Vilela CL. Technical note: Antimicrobial susceptibility of Portuguese isolates of *Staphylococcus aureus* and *Staphylococcus epidermidis* in subclinical bovine mastitis. J Dairy Sci. 2007; 90: 3242-6.
- 69. Mašlanková J, Pilipčincová I, Tkáčiková L. Phenotyping and genotyping of *Staphylococcus aureus* isolates of sheep origin. Acta Vet Brno. 2009; 78: 345-52.
- Huber H, Koller S, Giezendanner N, Stephan R, Zweifel C. Prevalence and characteristics of meticillinresistant *Staphylococcus aureus* in humans in contact with farm animals, in livestock, and in food of animal origin, Switzerland, 2009. Euro Surveill. 2010; 15: 19542.
- Friedrich A, Rau J, Horlacher S, Spohr M. Prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) in milk from dairy farms in Northern Württemberg. Tierarztliche Umschau. 2011; 66: 195-200.
- 72. Kreausukon K, Fetsch A, Kraushaar B, Alt K, Müller K, Krömker V, et al. Prevalence, antimicrobial resistance, and molecular characterization of methicillinresistant *Staphylococcus aureus* from bulk tank milk of dairy herds. J Dairy Sci. 2012; 95: 4382-8.
- 73. Fessler AT, Olde Riekerink, RGM, Rothkamp A, Kadlec K, Sampimon OC, Lam TJGM, et al. Characterization of methicillin-resistant *Staphylococcus aureus* CC398 obtained from humans and animals on dairy farms. Vet Microbiol. 2012; 160: 77-84.
- 74. Erskine RJ, Walker RD, Bolin CA, Bartlett PC, White DG. Trends in antibacterial susceptibility of mastitis pathogens during a seven-year period. J Dairy Sci. 2002; 85: 1111-8.
- 75. Makovec JA, Ruegg PL. Antimicrobial resistance of bacteria isolated from dairy cow milk samples submitted for bacterial culture: 8,905 samples (1994-2001). J Am Vet Med Assoc. 2003; 222: 1582-9.

- 76. Anderson KL, Lyman RL, Bodeis-Jones SM, White DG. Genetic diversity and antimicrobial susceptibility profiles among mastitis-causing *Staphylococcus aureus* isolated from bovine milk samples. Am J Vet Res. 2006; 67: 1185-91.
- Virgin JE, Van Slyke TM, Lombard JE, Zadoks RN. Short communication: methicillin-resistant *Staphylococcus aureus* detection in US bulk tank milk. J Dairy Sci. 2009; 92: 4988-91.
- D'amico DJ, Donnelly CW. Characterization of *Staphylococcus aureus* strains isolated from raw milk utilized in small-scale artisan cheese production. J Food Protect. 2011; 74: 1353-8.
- Coelho SMO, Reinoso E, Pereira IA, Soares LC, Demo M, Bogni C, et al. Virulence factors and antimicrobial resistance of *Staphylococcus aureus* isolated from bovine mastitis in Rio de Janeiro. Pesq Vet Bras. 2009; 29: 369-74.
- Saini V, McClure JT, Léger D, Keefe GP, Scholl, DT, Morck DW, et al. Antimicrobial resistance profiles of common mastitis pathogens on Canadian dairy farms. J Dairy Sci. 2012; 95: 4319-32.
- 81. Hata E, Katsuda K, Kobayashi H, Uchida I, Tanaka K, Eguchi M. Genetic variation among *Staphylococcus aureus* strains from bovine milk and their relevance to methicillin-resistant isolates from humans. J Clin Microbiol. 2010; 48: 2130-9.
- 82. Farzana K, Shah SNH, Jabeen F. Antibiotic resistance pattern against various isolates of *Staphylococcus aureus* from raw milk samples. J Res Sci. 2004; 15: 145-51.
- 83. Lee JH. Methicillin (Oxacillin)-resistant *Staphylococcus aureus* strains isolated from major food animals and their potential transmission to humans. Appl Environ Microbiol. 2003; 69: 6489-94.
- 84. Nam HM, Lee AL, Jung SC, Kim MN, Jang GC, Wee SH, et al. Antimicrobial susceptibility of *Staphylococcus aureus* and characterization of methicillin-resistant *Staphylococcus aureus* isolated from bovine mastitis in Korea. Foodborne Pathog Dis. 2011; 8: 231-8.
- 85. Moon JS, Lee AR, Kang HM, Lee ES, Kim MN, Paik YH, et al. Phenotypic and genetic antibiogram of methicillin-resistant staphylococci isolated from bovine mastitis in Korea. J Dairy Sci. 2007; 90: 1176-85.
- 86. Turutoglu H, Ercelik S, Ozturk D. Antibiotic resistance of *Staphylococcus aureus* and coagulase-negative staphylococci isolated from bovine mastitis. BullVet Res Inst Pulawy. 2006; 50: 41.
- Ünal N, Askar Ş, Macun HC, Sakarya F, Altun B, Yıldırım M. Panton–Valentine leukocidin and some exotoxins of *Staphylococcus aureus* and antimicrobial susceptibility profiles of staphylococci isolated from milks of small ruminants. Trop Anim Health Prod. 2012; 44: 573-9.
- Kumar R, Yadav BR, Singh RS. Genetic determinants of antibiotic resistance in *Staphylococcus aureus* isolates from milk of mastitic crossbred cattle. Cur Microbiol. 2010; 60: 379-86.
- 89. Kumar R, Yadav BR, Singh RS. Antibiotic resistance and pathogenicity factors in *Staphylococcus aureus* isolated from mastitic Sahiwal cattle. J Biosci. 2011; 36: 175-88.

- Intrakamhaeng M, Komutarin T, Pimpukdee K, Aengwanich W. Incidence of enterotoxin-producing MRSA in bovine mastitis cases, bulk milk tanks and processing Plants in Thailand. J Anim Vet Adv. 2012; 11: 655-61.
- Alian F, Rahimi E, Shakerian A, Momtaz H, Riahi M, Momeni M. Antimicrobial resistance of *Staphylococcus aureus* isolated from bovine, sheep and goat raw milk. Global Veterinaria. 2012; 8: 111-4.
- 92. Ateba CN, Mbewe M, Moneoang MS, Bezuidenhout CC. Antibiotic-resistant *Staphylococcus aureus* isolated from milk in the Mafikeng Area, North West province, South Africa. S Afr J Sci. 2010; 106: 1-6.
- Daka D, Solomon G, Yihdego D. Antibiotic-resistance Staphylococcus aureus isolated from cow's milk in the Hawassa area, South Ethiopia. Ann Clin Microbiol Antimicrob. 2012; 11: 26-37.
- 94. Suleiman AB, Umoh VJ, Kwaga JKP, Shaibu SJ. Prevalence and antibiotic resistance profiles of Methicillin resistant *Staphylococcus aureus* (MRSA) isolated from bovine mastitic milk in Plateau State, Nigeria. Intern Res J Microbiol. 2012; 2: 264-70.
- 95. Ferreira JP, Anderson KL, Correa MT, Lyman R, Ruffin F, Reller LB, et al. Transmission of MRSA between companion animals and infected human patients presenting to outpatient medical care facilities. PLoS One. 2011; 6: e26978.
- 96. Leonard FC, Markey BK. Meticillin-resistant *Staphylococcus aureus* in animals: a review. Vet J. 2008; 175(1): 27-36.
- 97. EFSA (European Food Safety Authority). Scientific Report of EFSA: Technical specifications on the harmonised monitoring and reporting of antimicrobial resistance in methicillin-resistant *Staphylococcus aureus* in food-producing animals and food. EFSA J. 2012; 10: 2897.
- 98. Kawada M, Okuzumi K, Hitomi S Sugishita C. Transmission of *Staphylococcus aureus* between healthy, lactating mothers and their infants by breastfeeding. J Hum Lact. 2003; 19: 411-7.
- 99. Peacock SJ, de Silva I, Lowy FD. What determines nasal carriage of *Staphylococcus aureus*? Trends Microbiol. 2001; 9: 605-10.
- 100. Rahmaniar RP, Yunita MN, Effendi MH, Yanestria SM. Encoding Gene for Methicillin Resistant Staphylococcus aureus (MRSA) Isolated from Nasal Swab of Dogs. Indian Veterinary Journal. 2020; **97** (02): 37 - 40.
- 101. Decline V, Effendi MH, Rahmaniar RP, Yanestria SM, Harijani N. Profile of antibiotic-resistant and presence of methicillin-resistant *Staphylococcus aureus* from nasal swab of dogs from several animal clinics in Surabaya, Indonesia, *Int. J. One Health*, 2020; 6(1): 90-94
- 102. Duquette RA, Nuttall TJ. Methicillin-resistant *Staphylococcus aureus* in dogs and cats: an emerging problem? J Small Anim Pract. 2004; 45: 591-7.
- 103. Catry B, Van Duijkeren E, Pomba MC, Greko C, Moreno MA, Pyörälä S, et al. Reflection paper on MRSA in foodproducing and companion animals: epidemiology and control options for human and animal health. Epidemiol Infect. 2010; 138: 626-44.
- 104. Cuny C, Friedrich A, Kozytska S, Layer F, Nübel U, Ohlsen K, et al. Emergence of methicillin-resistant *Staphylococcus aureus* (MRSA) in different animal species. Int J Med Microbiol. 2010; 300: 109-17.

- 105. Tyasningsih, W., Effendi, M. H., Budiarto, B., and Syahputra, I. R. Antibiotic Resistance to Staphylococcus aureus and Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated from Dairy Farms in Surabaya, Indonesia. *Indian Veterinary Journal*. 2019; **96** (11): 27-31.
- 106. Ramandinianto, S.C., Khairullah, A.R., Effendi, M.H. *MecA* gene and methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from dairy farms in East Java, Indonesia. BIODIVERSITAS, 2020; 21(8): 3562-3568.
- 107. Yunita, M.N., Effendi, MH, Rahmaniar, RP, Arifah S. and Yanestria. SM. Identification of SPA Gene for Strain Typing Of Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated From Nasal Swab of Dogs. *Biochem. Cell. Arch.* 2020; 20 (1): 2999-3004.
- 108. Sung JM, Lloyd DH, Lindsay JA. *Staphylococcus aureus* host specificity: comparative genomics of human versus animal isolates by multi-strain microarray. Microbiology. 2008; 154: 1949-59.
- 109. Khairullah, A. R., Ramandinianto, S.C., Effendi, M.H. A Review of Livestock-Associated Methicillin-Resistant Staphylococcus aureus (LA-MRSA) on Bovine Mastitis. *Sys Rev Pharm 2020;11(7):172-183.*
- 110. Ramandinianto, S.C., Khairullah, A.R., Effendi, M.H., Tyasningsih, W. and Rahmahani, J. Detection of Enterotoxin type B gene on Methicillin Resistant *Staphylococcus aureus* (MRSA) isolated from raw milk in East Java, Indonesia. *Sys Rev Pharm* 2020;11(7):290-298.
- 111. van Rijen MM, Van Keulen PH, Kluytmans JA. Increase in a Dutch hospital of methicillin-resistant *Staphylococcus aureus* related to animal farming. Clin Infect Dis. 2008; 46: 261-3.
- 112. Cuny C, Nathaus R, Layer F, Strommenger B, Altmann D, Witte W. Nasal colonization of humans with methicillin-resistant *Staphylococcus aureus* (MRSA) CC398 with and without exposure to pigs. PLoS One. 2009; 27(4): e6800.
- 113. Cefai C, Ashurst S, Owens C. Human carriage of methicillin-resistant *Staphylococcus aureus* linked with pet dog. Lancet. 1994; 344: 539-40.
- 114. Manian FA. Asymptomatic nasal carriage of mupirocin-resistant, methicillin-resistant *Staphylococcus aureus* (MRSA) in a pet dog associated with MRSA infection in household contacts. Clin Infect Dis. 2003; 36: 26-28.
- 115. Baptiste KE, Williams K, Willams NJ, Wattret A, Clegg PD. Methicillin-resistant staphylococci in companion animals. Emerg Infect Dis. 2005; 11: 1942-4.
- 116. van Duijkeren E, Wolfhagen MJ, Heck ME, Wannet WJ. Transmission of a panton-valentine leucocidinpositive, methicillin-resistant *Staphylococcus aureus* strain between humans and a dog. J Clin Microbiol. 2005; 43: 6209-11.
- 117. Faires M, Tater K, Weese JS. An investigation of methicillin-resistant *Staphylococcus aureus* colonization in people and pets in the same household with an infected person or infected pet. J Am Vet Med Assoc. 2009; 235: 540.
- 118.Weese JS, Avery BP, Reid-Smith RJ. Detection and quantification of methicillin-resistant *Staphylococcus aureus* (MRSA) clones in retail meat products. Appl Microbiol. 2010; 51(3): 338-42.

- 119. Ferry T, Bes M, Dauwalder O, Meugnier H, Lina G, Forey F, et al. Toxin gene content of the Lyon methicillin-resistant *Staphylococcus aureus* clone compared with other pandemic clones. J Clin Microbiol. 2006; 44: 2642-4.
- 120. Tristan A, Ferry T, Durand G, Dauwalder O, Bes M, Lina G, et al. Virulence determinants in community and hospital meticillin-resistant *Staphylococcus aureus*. J Hosp Infect. 2007; 65: 105-9.
- 121.EFSA (European Food Safety Authority). Foodborne antimicrobial resistance as a biological hazard Scientific Opinion of the Panel on Biological Hazards Public consultation (Question No EFSA-Q-2007-089). EFSA J. 2008; 765: 2-87.
- 122. Graham PL, Lin SX, Larson EL. A U.S. population-based survey of *Staphylococcus aureus* colonization. Ann Intern Med. 2006; 144(5): 318-25.
- 123. Smith TC, Gebreyes WA, Abley MJ, Harper AL, Forshey BM, Male MJ, et al. Methicillin-resistant *Staphylococcus aureus* in pigs and farm workers on conventional and antibiotic-free swine farms in the USA. PLoS ONE. 2013; 8: e63704.
- 124. Wertheim HFL, Melles DC, Vos MC, van Leeuwen W, van Belkum A, Verbrugh HA, et al. The role of nasal carriage in *Staphylococcus aureus* infections. Lancet Infect Dis. 2005; 5(12): 751-62.
- 125. Fritz SA, Epplin EK, Garbutt J, Storch GA. Skin infection in children colonized with community-associated methicillin-resistant *Staphylococcus aureus*. J Infect. 2009; 59(6): 394-401.
- 126. Murphy BP, O'Mahony E, Buckley JF, O'Brien S, Fanning S. Characterization of *Staphylococcus aureus* isolated from dairy animals in Ireland. Zoonoses Public Health. 2010; 57(4): 249-57.
- 127. Gunaydin B, Aslantas O, Demir C. Detection of superantigenic toxin genes in *Staphylococcus aureus* strains from subclinical bovine mastitis. Trop Anim Health Pro. 2011; 43(8): 1633-7.
- 128. Neder VE, Canavesio VR, Calvinho LF. Presence of enterotoxigenic *Staphylococcus aureus* in bulk tank milk from Argentine dairy farms. Rev Argent Microbiol. 2011; 43(2): 104-6.
- 129. Arcuri EF, Angelo FF, Guimaraes MFM, Talon R, Borges MF, Leroy S, et al. Toxigenic status of *Staphylococcus aureus* isolated from bovine raw milk and Minas frescal cheese in Brazil. J Food Prot. 2010; 73(12): 2225-31.
- 130. Scherrer D, Corti S, Muehlherr JE, Zweifel C, Stephan R. Phenotypic and genotypic characteristics of *Staphylococcus aureus* isolates from raw bulk-tank milk samples of goats and sheep. Vet Microbiol. 2004; 101(2): 101-7.
- 131. Peles F, Wagner M, Varga L, Hein I, Rieck P, Gutser K, et al. Characterization of *Staphylococcus aureus* strains isolated from bovine milk in Hungary. Int J Food Microbiol. 2007; 118(2): 186-93.
- 132.do Carmo LS, Dias RS, Linardi VR, de Sena MJ, dos Santos DA, de Faria ME, et al. Food poisoning due to enterotoxigenic strains of Staphylococcus present in Minas cheese and raw milk in Brazil. Food Microbiol. 2002; 19(1): 9-14.
- 133. Le Loir Y, Baron F, Gautier M. *Staphylococcus aureus* and food poisoning. Genet Mol Res. 2003; 2: 63-76.
- 134.Kowalski RP, Karenchak LM, Romanowski EG. Infectious disease: changing antibiotic susceptibility. Ophthalmol Clin. 2003; 16:1-9.

- 135.Lowy FD. Staphylococcus aureus infections. N Engl J Med. 1998; 339(8): 520-32.
- 136. Makgotlho PE, Kok MM, Hoosen A, Lekalakala R, Omar S, Dove M. Molecular identification and genotyping of MRSA isolates. FEMS Imunol Med Mic. 2009; 57(2): 104-15.
- 137. Michel M, Gutmann L. Methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant enterococci: therapeutic realities and possibilities. Lancet. 1997; 349(9069): 1901-6.
- 138. Birmingham MC, Rayner CR, Meagher AK, Flavin SM, Batts DH, Schentag JJ. Linezolid for the treatment of multidrug-resistant, gram-positive infections: experience of a compassionate-use program. Clin Infect Dis. 2003; 36(2): 159-68.
- 139. Furtado S, Bhat RM, Rekha B, Sukumar D, Kamath GH, Martis J, et al. The clinical spectrum and antibiotic sensitivity patterns of staphylococcal pyodermas in the community and hospital. Indian J Dermatol. 2014; 59(2): 143-50.
- 140.Wang R, Braughton KR, Kretschmer D, Bach THL, Queck SY, Li M. Identification of novel cytolytic peptides as key virulence determinants for community-associated MRSA. Nat Med. 2007; 13(12): 1510-4.
- 141.Todd ECD, Greig JD, Bartleson CA, Michaels BS. Outbreaks where food workers have been implicated in the spread of foodborne disease. J Food Prot. 2008; 71: 2582-95.
- 142. Bystron J, Podkowik M, Korzekwa K, Lis E, Molenda J, Bania J. Characterization of borderline oxacillinresistant *Staphylococcus aureus* isolated from food of animal origin. J Food Prot. 2010; 73: 1325-7.
- 143. Valero A, Perez-Rodriguez F, Carrasco E, Fuentes-Alventosa JM, Garcia-Gimeno RM, Zurera G. Modelling the growth boundaries of *Staphylococcus aureus*: effect of temperature, pH and water activity. Int J Food Microbiol. 2009; 133: 186-94.
- 144.Oliver SP, Murinda SE, Jayarao BM. Impact of antibiotic use in adult dairy cows on antimicrobial resistance of veterinary and human pathogens: A comprehensive review. Foodborne Pathog Dis. 2011; 8(3): 337-55.
- 145. Effendi M.H., Hisyam M.A.M., Hastutiek P., and Tyasningsih W. Detection of coagulase gene in *Staphylococcus aureus* from several dairy farms in East Java, Indonesia, by polymerase chain reaction, *Vet. World*, 2019; **12**(1): 68-71.
- 146. Effendi, M.H., Oktavianto, A and Hastutiek, P. Tetracycline Resistance Gene in Streptococcus Agalactiae Isolated from Bovine Subclinical Mastitis in Surabaya, Indonesia. *Philipp. Journal of Veterinary Medicine*. 2018; **55** (SI): 115-120.
- 147. Dhanashekar R, Akkinepalli S, Nellutla A. Milk-borne infections. An analysis of their potential effect on the milk industry. GERMS. 2012; 2(3): 101-9.