

A Study on Urinary Tract Infection and Metabolic Diseases in Stone Patients

Abdul Tawab Zia¹, Muhammad Qamar uz Zaman¹, Sajjid Abbas²

¹Department of general practice, Rural Health Center, Midh ranjha, tehsile Kotmomin, Sargodha, Pakistan

²DHQ Teaching hospital, Sargodha, Pakistan

Article History:

Submitted: 12.04.2021

Accepted: 19.05.2021

Published: 26.05.2021

ABSTRACT

Background and objectives: The main objective of the study is to analyze the Urinary Tract Infection and metabolic diseases in Stone Patients among local population of Pakistan.

Material and methods: This cross sectional study was conducted at DHQ teaching hospital, Sargodha during June 2019 to June 2020. The definite history of the multitude of patients were accumulated and 24 hour pee test was gathered from every patient and sent for PH, explicit gravity, Creatinine, uric corrosive, calcium, phosphate, oxalate, citrate and magnesium. 24-hour pee tests were gathered in plastic boxes, which don't respond artificially by standard techniques, and were put away at 2°C-8°C. Likewise, blood test of every patient was additionally sent for serum levels of urea, creatinine, uric corrosive, phosphate and calcium.

Results: The data were collected from 100 patients with the mean age 38 ± 7.75 years. There were 35 male

and 65 female patients who were selected this investigation. The primary introducing grievance was amble torment on the influenced side for example in 79.0% patients, trailed by haematuria and consuming micturition. Dominant part of the patients for example 94.0%, were analysed as having renal stone or ureteric stone.

Conclusion: It is concluded that recurrence of metabolic variations from the norm is exceptionally high in patients with urolithiasis and hyperoxaluria, hypercalciuria and hypocitraturia are the main metabolic irregularities saw in these patients. Stone sickness is an expanding and significant general health issue with high recurrence of bladder stone.

Keywords: Urinary, Tract, Infection, Patients, Metabolic

Correspondence:

Department of general practice, Rural Health Center, Midh ranjha, tehsile Kotmomin, Sargodha, Pakistan, E-mail: tawabgujar107@gmail.com

INTRODUCTION

Urolithiasis is a disease which is common now a days in the whole world. It is assessed that roughly 2% of the populace experience renal stone illness at times in their existence with top rate in second and third many years of life. There are a few sorts of urinary stones, and they are ordered by compound composition (Taylor EN, *et al.*, 2005). Calcium oxalate is the significant segment of by far most of stones. A few components, for example, age, sexual orientation, atmosphere, metabolic variations from the norm and heredity are related to the improvement of urinary stones. Metabolic variations from the norm are the main elements since they can be altered to forestall the danger of urinary stones (Courbebaisse M, *et al.*, 2017).

Inhabiting urinary catheters are standard clinical gadgets used in both clinic and nursing home settings to mitigate urinary maintenance and urinary incontinence (Moe OW, 2006). Of the very nearly 100 million catheters that are sold yearly around the world, one-fourth of them are sold in the United States (Robertson WG, *et al.*, 1979). The most widely recognized urinary catheter being used is the Foley inhabiting urethral catheter, a shut sterile framework that is included a cylinder embedded through the urethra and held set up by an inflatable to permit urinary seepage of the bladder (Joseph KC, *et al.*, 2005). In spite of the fact that these gadgets were initially intended for transient use in patients, inhabiting catheter use is currently typical in the long haul setting (Chhiber N, *et al.*, 2014). Infections of the urinary tract related with catheter use are huge not just due their high frequency and ensuing financial expense yet additionally on account of the serious sequelae that can result (Chaudhary A, *et al.*, 2010).

Patients with confounded UTI should be treated by compelling antimicrobial treatment just as suitable urological medication to eliminate inclining factors when the manifestations

are related, for example, micturition torment, dysuria, urinary recurrence, and low or high fever (Moro F, *et al.*, 2005). Paradoxically, Asymptomatic Bacteriuria (ASB) is recognized by segregation of a predetermined quantifiable measure of microorganisms in a properly gathered pee example acquired from a patient without indications or signs referable to urinary infection (Aggarwal KP, *et al.*, 2013).

Aims and objectives

The main objective of the study is to analyse the Urinary Tract Infection and metabolic diseases in stone patients among local population of Pakistan.

METHODOLOGY OF THE STUDY

This cross sectional study was conducted at DHQ teaching hospital, Sargodha during June 2019 to June 2020. There were all out 100 patients of the two sexual orientations who were chosen for this examination. The data was gathered through an arbitrary testing strategy method.

Inclusion criteria

Patients with one or the other various or repetitive urolithiasis were remembered for this investigation.

Exclusion criteria

- Patients with some other sickness were prohibited from this examination.
- Patients with ongoing renal disappointment, persistent liver sickness and with history of any constant medication use were rejected.

Data collection

The data were collected from 100 patients. The positive history of the large number of patients were collected and 24 hour pee test was assembled from each patient and sent for PH, express grav-

ity, creatinine, uric destructive, calcium, phosphate, oxalate, citrate and magnesium. 24-hour pee tests were accumulated in plastic boxes, which don't react misleadingly by standard procedures, and were taken care of at 2°C-8°C. Moreover, blood trial of each patient was furthermore sent for serum levels of urea, creatinine, uric destructive, phosphate and calcium. The serum levels of metabolic limits were assessed by standard compound strategy. All patients by then had legitimate method after fulfilment of all workup and stones were transported off pathology research office for compound assessment to consider the stone structure.

The data were collected and analysed through SPSS (Version 21.0). All the values were expressed in mean and standard deviation.

RESULTS

The data were collected from 100 patients with the mean age 38 ± 7.75 years. There were 35 male and 65 female patients who were selected this investigation. The primary introducing grievance was amble torment on the influenced side for example in 79.0% patients, trailed by haematuria and consuming micturition. Dominant part of the patients for example 94.0%, were analysed as having renal stone or ureteric stone. Just 38.0% patients gave intermittent stones while staying 62.0% had stone unexpectedly. Substance examination of stones after complete strategy had demonstrated calcium oxalate stone in 82.5% patients (Table 1). Hyperoxaluria was the most normally noticed metabolic anomaly and was found in 12 patients. Other huge metabolic variations from the norm were hypercalciuria, Hypercalcemia, hypocitraturia and hyperuricemia (Table 2).

Table 1: Descriptive statistics for different variables

Features	%Age
Presenting complaint	
Lumber pain	79
Hematuria	13
Burning micturition	8
Diagnosis	
Renal stone	63
Ureteric stone	21
Renal+ureteric stone	10.06.0
Urinary bladder stone	
Recurrent stone:	
Yes	38
No	62
Family history of urolithiasis	
Yes	64
No	36
Stone composition on stone analysis	
Calcium oxalate	82.5
Calcium phosphate	2.5
Uric acid	11.5
Struvite	1.5
Cystine	2

Table 2: Frequency of metabolic diseases in selected participants

Metabolic abnormality	Frequency	%Age
Hyperoxaluria	12	12.1
Hypercalciuria	17	16.9
Hypocitraturia	11	11.2
Hypernatruria	19	18.8
Hyperuricosuria	3	3
Hypomagnesuria	8	7.9

Hyperphosphaturia	13	13.1
Hypercalcemia	5	5

DISCUSSION

Urinary stones in its various structures are the third most basic difficulty of the urinary tract. Math illness is the commonest urological sickness in Pakistan. It has been evident for quite a long while that the occurrence paces of lithiasis shift significantly, from mainland to land-mass as well as between neighbouring areas of a nation, regardless of whether one takes into consideration contrasts in technique and models determination among the study of disease transmission studies (Aggarwal KP, *et al.*, 2013). The lifetime pervasiveness of urinary stones has expanded all through the twentieth century and happens in up to 15% of the populace. It is commonly acknowledged that stones happen more ordinarily in guys than females (Butterweck V and Khan SR, 2009). Our discoveries certify with this sex contrast as announced by others. No age bunch is saved to urinary stone illness in Pakistan however an adjustment in the age example of patients of urolithiasis has been accounted for in industrialized countries (Daudon M, *et al.*, 2004). In our examination, the primary introducing grumbling was blunder torment for example in 79.0% patients. Elfadil GA *et al.* had likewise discovered flank torment as the head introducing grievance in his investigation for example in 67% patients (Cachat F, *et al.*, 2004). The consequences of our investigation have demonstrated a solid hereditary inclination to urinary stone infection as 64.0% patients had family background of urolithiasis. This hereditary factor is additionally upheld by investigations of Kirac M *et al.* and Majalan NN *et al.* who had discovered a positive family ancestry in 67.0% and 53.1% patient's respectively (Daudon M, *et al.*, 2004). Then again, Elfadil GA *et al.* (Chaudhary A, *et al.*, 2010) had discovered this in just 20% of their patients. We had additionally discovered 38.0% patients with intermittent urinary stones and the significant stone segment was calcium oxalate in our examination which was likewise found by Elfadil GA *et al.* (Chaudhary A, *et al.*, 2010). But in an investigation by Androulakakis *et al.*, the primary segments of urinary stones in Europe, in diminishing request, are struvite, calcium phosphate and calcium oxalate (Bakane BC, *et al.*, 1999).

In our investigation, metabolic anomalies were found in 90.5% patients, while there was no metabolic variation from the norm in just 9.5% patients which is a lot of practically identical to numerous past examinations. In an investigation by Amaro *et al.*, 62.2% of patients had various metabolic variations from the norm; in any case, the patients didn't have repetitive calcium oxalate stones (Yoshida O, *et al.*, 1999). Accordingly, it very well may be assumed that numerous metabolic anomalies are more normal in patients with intermittent calcium oxalate stones (Sternberg K, *et al.*, 2005).

CONCLUSION

It is concluded that recurrence of metabolic variations from the norm is exceptionally high in patients with urolithiasis and hyperoxaluria, hypercalciuria and hypocitraturia are the main metabolic irregularities saw in these patients. Stone sickness is an expanding and significant general health issue with high recurrence of bladder stone.

REFERENCES

1. Taylor EN, Stampfer MJ, Curhan GC. Obesity, weight gain and the risk of kidney stones. *Journal of the American Medical Association.* 2005; 293(4): 455-462.
2. Courbebaisse M, Bertoye CP, Bertocchio JP, Baron S, Maruani G, Briand S, *et al.* Nephrolithiasis of adult: From mechanisms to preventive medical treatment. *Revue Medicale Internationale.* 2017; 38(1): 44-52.

3. Moe OW. Kidney stones: Pathophysiology and medical management. *The Lancet*. 2006; 367(9507): 333-344.
4. Robertson WG, Heyburn PJ, Peacock M, Hanes FA, Swaminathan R. The effect of high animal protein intake on the risk of calcium stone-formation in the urinary tract. *Clin Sci*. 1979; 57(3): 285-288.
5. Joseph KC, Bharat B, Parek H, Joshi MJ. Inhibition of growth of urinary type calcium hydrogen phosphate dihydrate crystals by tartaric acid and tamarind. *Curr Sci*. 2005; 88: 1232-1238.
6. Chhiber N, Sharma M, Kaur T, Singla SK. Mineralization in health and mechanism of kidney stone formation. *Int J Pharm Sci Invent*. 2014; 3(10): 25-31.
7. Chaudhary A, Singla SK, Tandon C. In vitro evaluation of Terminalia arjuna on calcium phosphate and calcium oxalate crystallization. *Indian J Pharm Sci*. 2010; 72(3): 340-345.
8. Moro F, Mancini M, Tavolini IM, De Marco V, Bassi P. Cellular and molecular gateways to urolithiasis: a new insight. *Urologia Internationalis*. 2005; 74: 193-197.
9. Aggarwal KP, Narula S, Kakkar M, Tandon C. Nephrolithiasis: Molecular mechanism of renal stone formation and the critical role played by modulators. *BioMed Research International*. 2013; 2013: 292953.
10. Butterweck V, Khan SR. Herbal medicines in the management of urolithiasis: Alternative or complementary. *Planta Medica*. 2009; 75(10): 1095-1103.
11. Daudon M, Bounxouei B, Santa Cruz F, Leite da Silva S, Diouf B, Angwafo FF, *et al*. Composition of renal stones currently observed in non-industrialized countries. *Prog Urol*. 2004; 14(6): 1151-1156.
12. Cachat F, Barbey F, Guignard JP. Urinary calculi epidemiology in children. *Rev Med Suisse Romande*. 2004; 124(8): 433-437.
13. Bakane BC, Nagtilak SB, Patil B. Urolithiasis: A tribal scenario. *Indian J Pediatr*. 1999; 66(6): 863-866.
14. Yoshida O, Terai A, Ohkawa T, Okada Y. National trend of the incidence of urolithiasis in Japan from 1965 to 1995. *Kidney Int*. 1999; 56(5):1899-1904.
15. Sternberg K, Greenfield SP, Williot P, Wan J. Pediatric stone disease: An evolving experience. *J Urol*. 2005; 174: 1711-1714.