

Circulatory Leptin Levels as a Key Link in the Potential Association between Insulin Resistance and Vitamin D Deficiency: A Review Article

Mahmoud Abu-Samak¹, Beisan A Mohammad², Ibrahim Mosleh³, Omar Abdel-Majeed¹, Abeer Braham¹, Shady H Awwad⁴, Ahamed Abu-Zaiton⁵, Abdel Rahman M. Tawaha⁶

¹ Department of Clinical Pharmacy and Therapeutics, Applied Science Private University, Amman, Jordan

² Pharm D Department, Fakeeh College for Medical Sciences, Jeddah, Kingdom of Saudi Arabia

³ Department of Clinical Laboratories Sciences, University of Jordan, Amman, Jordan

⁴ Department of Pharmaceutical Chemistry & Pharmacognosy, Applied Science Private University, Amman, Jordan

⁵ Department of Biological Sciences, Al al-bayt University, Al-Mafraq, Jordan

⁶ Department of Biological Sciences, Al Hussein bin Talal University, P.O. Box 20, Maan, Jordan

Correspondence: Mahmoud S Abu-Samak

Department of Clinical Pharmacy and Therapeutics, Applied Science Private University, Al Arab Street 21, Amman 11931, Jordan

Email m_abusamak@asu.edu.jo

ABSTRACT

Vitamin D deficiency (VDD) and diabetes mellitus (DM) are worldwide problems. Many recent studies have shown a high prevalence of T2DM and VDD in Mediterranean countries including Jordan. Remarkably, VDD is more prevalent in obese people with inverse association with 25-hydroxy vitamin D (25OHD). Nevertheless, randomized clinical trials (RCTs) point did not confirm that yet in particular the association with an obesity marker (leptin hormone). Leptin is involved in the insulin resistance (IR) pathogenesis and development of T2DM. Some RCTs showed that the treatment of VDD by 1,25OHD₂D₃ (VD₃) may improve the control of diabetes and insulin sensitivity or decrease the risk of disease. Conversely, leptin levels were positively associated with 25OHD levels. Overall, taking into account the U-shaped curve, it seems that the association between VD₃ supplementation is a dose dependent. In this context, it can be concluded that when VD intake is below, the serum leptin level will be low and vice versa. VD supplementation may cause raising serum leptin as shown in this review. Therefore, we suggest further clinical trials wither to confirm or negation of existence of diabetogenic or anti-diabetogenic effects for VD₃ supplementations.

Keywords: VDD, diabetes, T2DM, vitamin D, leptin, obesity, insulin resistance

Correspondence:

Mahmoud Abu-Samak

Department of Clinical Pharmacy and Therapeutics, Applied Science Private University, Al Arab Street 21, Amman 11931, Jordan
Email m_abusamak@asu.edu.jo

INTRODUCTION

Vitamin D deficiency (VDD), obesity and diabetes mellitus (DM) are worldwide problems. There has been growing interest in the association of type 2 DM (T2DM) with VDD in relation to obesity (Smith and Singleton, 2013). Many studies have shown a high prevalence of T2DM and VDD in many populations including Mediterranean countries (Qatatsheh *et al.*, 2015; Al-Shaer *et al.*, 2016). Some reports have shown that VDD was more prevalent in obese people (Vimaleswaran *et al.*, 2015; Hajimohammadi *et al.*, 2016; Chadt *et al.*, 2018). An inverse association between 25-hydroxy vitamin D (25OHD) levels and BMI was also observed (Ruiz-Ojeda *et al.*, 2018). In this manner, leptin hormone, the obesity gene product, is one of the most important obesity and insulin resistance markers (Yadav *et al.*, 2013). However, the association between VDD and serum leptin levels is not settled yet (Kim *et al.*, 2013; Hajimohammadi *et al.*, 2016). Meta-analysis of data from 6 RCTs did not find a significant change in plasma leptin concentrations after vitamin D₃ (VD₃) intervention (Dinca *et al.*, 2016). Contrary to Dinca *et al.* (2016), a significant positive association between 25OHD and leptin has been reviewed by Hajimohammadi *et al.* (2016). High prevalence of T2DM among people with VDD were spread dramatically during last decade. Therefore, the current review aims highlight the association between vitamin D deficiency and serum leptin as a potential key link and circulatory marker of insulin resistance.

Vitamin D Deficiency

Vitamin D (VD) is an essential fat-soluble vitamin, endogenously produced by the action of sunlight on 7-dehydrocholesterol in skin (also known as D₃ or cholecalciferol) or obtained from dietary food stuff as either VD₂ (known as ergocalciferol) or VD₃ (Hewison, 2012). VD occurs in food in small amounts, mainly in the raw material of an animal origin, whereas in plant products VD is not present. Chemically, the various forms of VD belong to secosteroids; a biochemical class of 'broken' ring steroids: VD₂, and VD₃ or cholecalciferol are the two major forms which are known collectively as VD or calciferol (Newberry *et al.*, 2014).

Circulatory levels of 25OHD as well as other calciotropic hormones and predictors such as parathyroid hormone (PTH), FGF23, calcitonin and sex hormones are involved in the interrelationships that maintain normal calcium levels (Kovacs, 2014; Mundy and Guise, 1999). The cutaneous synthesis of VD₃ depends on the age, skin color, latitude, season of the year and time of the day (Newberry *et al.*, 2014), sunscreen use, air pollution (Hosseini-nezhad and Holick, 2013) and it is also proportional to the skin surface area exposed to the sun (Wimalawansa, 2016). This way of VD production is considered safe since prolonged exposure to UVB rays will not produce toxic levels of VD₃ due to the conversion of pre VD₃ to inactive metabolites tachysterol and lumisterol (Zhang and Naughton, 2010). After formation VD₃ will be discharged from keratinocytes plasma membrane to the dermal capillary tube by DBP, then VD₃ will be assimilated into chylomicrons, released into

Circulatory Leptin Levels as a Key Link in the Potential Association between Insulin Resistance and Vitamin D Deficiency: A Review Article

lymphatic system and enter the blood (Hossein-nezhad and Holick, 2013). Cutaneous synthesis of VD may not be sufficient to produce adequate amounts of VD, because of that the VD supplementations are advisable to prevent VDD (Chen *et al.*, 2007). Exogenous dietary form of VD is absorbed in the jejunum and ileum with the simplifying of bile salts (Teske *et al.*, 2016). In healthy human, there is no difference between VD₂ and VD₃ in absorption rate, the efficiency of VD absorption varies between 55% and 99%. Patients with fat malabsorption, particularly steatorrhea are unable to absorb VD (Nair and Maseeh, 2012). The absorbed VD is incorporated into chylomicrons before entering the circulation. 85-90% of VD forms are transported by a liver protein DBP to the target tissues (Jovičić *et al.*, 2012). 10-15% of the total circulating VD is bound to albumin and less than 1% is in the free form (Powe *et al.*, 2013).

Vitamin D deficiency and health problems

VDD remains a public health concern, due to its association with many health outcomes, VDD is linked to increased risk of cancer, cardiovascular disease, diabetes, autoimmune disease (Lerchbaum and Obermayer, 2012), metabolic disorders, infectious diseases (Theodoratou *et al.*, 2014), and multiple sclerosis. The correlation between vitamin deficiencies including vitamins B12 and VD with obesity was previously mentioned (Abu-Samak *et al.*, 2008; Pearce and Cheetham, 2010). High prevalence of VDD may be related to several factors, such as less VD photosynthesis in response to insufficient ultra-violet radiation, low VD intake, age, obesity, hyperlipidemia dark-skinned, stress can also contribute to VDD (Mallah, *et al.*, 2011; Gonzalez, 2014; Abu-Samak *et al.*, 2019).

Type 2 diabetes mellitus (T2DM) and Insulin resistance

T2DM is a multifactorial metabolic disorder resulting from a complex inheritance-environment interaction along with other risk factors (Gudjinu and Sarfo, 2017). The risk factors in children are similar to those in adults: ethnicity, family history, hyperlipidemia, sedentary lifestyle, and obesity (Abu-Hasheesh *et al.*, 2010; Abu-Samak *et al.*, 2013; Abu-Samak *et al.*, 2018; Abu-Taha *et al.*, 2019). Severe obesity represented a major risk factor for the development of T2DM (Chadt *et al.*, 2018). Insulin resistance (IR) is the key factor in linking between obesity and T2DM. T2DM accounts the majority of diabetic cases (Gao *et al.*, 2017). In many countries, T2DM became a significant health problem due to an increase in obesity among the young, unhealthy dietary habits and a sedentary lifestyle (Jarab *et al.*, 2018). It is well known that obesity increases the risk for T2DM through induction of IR (Abu-Hasheesh *et al.*, 2010). Obesity is the excessive or abnormal accumulation of fat in the body sufficient to increase overall morbidity and mortality. The importance of obesity in the etiology of T2DM is highlighted by the fact that the development of diabetes is due to rising rates of obesity (most of T2DM are obese). Moreover, weight loss in obese patients of T2DM can ameliorate or even terminate the disorder. Before the onset of clinical diabetes, during the early stages of T2DM pathogenesis, also known as prediabetics, IR continues to be questioned so it is characterized by high insulin levels. This explains the hypothesis that IR could be the primary reason, resulting in a compensatory increase in insulin secretion that ultimately cannot be maintained by the exhausted pancreas which cannot keep up with insulin demands (Tong *et al.*, 2016).

Insulin Resistance: Role of leptin

Adipose tissue is the primary source for IR mediators (González, 2012). Lipotoxicity (excess free fatty acids) that decreases skeletal muscle insulin sensitivity by interfering with insulin receptor substrate -1 (IRS-1) signaling and central adiposity that increases IR appears to include dysregulated secretion of leptin which is produced the fat tissue. Leptin is 167-amino-acid metabolic peptide, discovered in 1991 (Ferguson, 2014) that controls and maintains body weight by regulating appetite and fat metabolism. Leptin hormone acts centrally to control satiety and enhance insulin sensitivity (Lanzerstorfer *et al.*, 2015). Obesity can be a result of leptin resistance. Therefore leptin resistance is likely to be involved in the development of T2DM (Finucane *et al.*, 2009). Leptin deficiency is also considered to be noteworthy in the pathogenesis of IR in uncontrolled T2DM. Accordingly, many studies have considered that leptin has anti-diabetogenic effects via improving IR or by mediating the release of insulin from pancreatic β cells (D'Souza *et al.*, 2014). Animal models also, showed that leptin administration reversed diabetes in lipoatrophic mice. Finally, diet and leptin treatment should be thoroughly explored as a method of diabetes control (Silva *et al.*, 2017).

The association between VDD and Insulin Resistance : Leptin role

Adipose tissue is the primary source of mediators of IR (Frayn, 2001). The mechanisms by which fat tissue, particularly central (abdominal) adiposity, increases IR continues to be explained and appears to include: lipotoxicity which decrease skeletal muscle insulin sensitivity by interfering with IRS signaling; and dysregulated secretion of the anti-diabetogenic hormone, leptin in the fat tissue which acts centrally to control satiety and enhance insulin sensitivity (D'Souza *et al.*, 2014). IR in turn is strongly associated with increased leptin hypersecretion which may play the key role in the development of insulin insensitivity (Klötting and Blüher, 2014). Parental history of diabetes has been linked also as a potential predictor (Abu-Hasheesh *et al.*, 2010). In concerning to VDD, numerous global studies have reported a negative correlation between BMI and serum 25OHD levels (Konradsen *et al.*, 2008; Lagunova *et al.*, 2009; Khawaja *et al.*, 2017). The lack of morning sun exposure was one of the predominate percentages in this study as previously agreed (Mead, 2008). Some of reports have reviewed that VDD increases DM risk (Boucher *et al.*, 2004; Pittas *et al.*, 2019) referred that VDD enhances IR and develops T2DM. A correlation study by (Gandhe, 2013) observed a negative association between 25OHD levels and the risk of DM. Further, recent RCTs showed that treatment of VDD by VD₃ may improve the control of diabetes in Asian subjects (Upreti *et al.*, 2018). Accordingly, it has concluded that the effect of VD supplementation might play a role in the prevention of T1DM (Monnier and Colette, 2010). Recently, in prediabetic patients with VDD, high dose of VD₃ improved insulin sensitivity and decreases risk of progression toward diabetes (Niroomand *et al.*, 2019). However, the association of VDD with DM is still not clarified and needs more studies particularly clinical trials. Therefore, vitamin D receptors (VDR) in obese subjects has been drawing attention in related VD studies including it is a potential mediating role in the effect of VD on insulin hormone action (Knekt *et al.*, 2008). Despite that Kim *et al.* (2013) have mentioned no correlation between BMI and 25(OH)D, many studies had shown that the VDD is more prevalent in obese people with inverse association

Circulatory Leptin Levels as a Key Link in the Potential Association between Insulin Resistance and Vitamin D Deficiency: A Review Article

between 25OHD levels and BMI (Eganet *et al.*,2014). These observations confirmed an existence of relationship between serum leptin and 25OHD in men and women with VDD. Nevertheless, conflicting results have been reported by Kim *et al.*(2013). For example, circulatory 25OHD variation has not been associated with BMI values (Forsythe *et al.*, 2011). Conversely, an inverse correlation has been showed between serum 25OHD levels with fat volume (Vimaleswaran *et al.*,2015) and with leptin in different age groups (Hajimohammadi *et al.*,2016). In relation to VDD, it was noted that each increase in BMI unit corresponds to a 1.15% reduction in the 25OHD level (Harroud and Richards, 2018). Similarly, but in relation to fat weight, a 1% increase in fat weight was associated with a (1.15 ± 0.55) nmol/L reduction in serum 25OHD level (Lenders *et al.*,2009).

On the other hand, findings of randomized clinical trials (RCTs) that examined the effect of different doses of VD₃ supplementations on serum leptin levels in healthy subjects or patients, were also mixed. In healthy subjects, no significant change in circulatory leptin levels had been noted according to different protocols of VD₃ doses (Dinca *et al.*,2016; Duggan *et al.*, 2015; Mousa *et al.*, 2019). Conversely, significant increase (Hajimohammadi *et al.*,2016.) or decrease (Vahdat *et al.*,2016) in the mean value of leptin levels were observed. It seems that high doses of VD₃ have significant influence to decrease of leptin levels in some populations. It has shown that one shoot of a very high dose (600,000 IU D₃) significantly reduced leptin levels in obese subjects (Mai *et al.*, 2017). Also, high doses of VD₃ (50,000 IU/w for 6 weeks) also reduced significantly the mean of BMI and body weight in Iranian population (Entezari *et al.*,2018). BMI variation based on the mean age value might be positively correlated with body fat mass (Mott *et al.*, 1999). Furthermore, an inverse correlation between serum 25OHD levels with fat volume was more significant than BMI (Vimaleswaran *et al.*, 2015). This is because of leptin is more associated with obesity than BMI as mentioned by Abu-Samak *et al.*(2011) or due to the absorption of vitamin D by adipose tissue (Wortzman *et al.* 2000), which clarifies, in part, the inverse association between 25OHD and obesity. Although observational studies contributed contraindicating findings that to high heterogeneity sources of heterogeneity such as age and health condition, it seems that the effect of VD₃ on blood leptin levels is a dose dependent (Hajimohammadi *et al.*,2016).

Leptin hypersecretion by adipocyte hypertrophy may plays the key role in the development of insulin insensitivity in obese people (Targher *et al.*, 2007) as well as may impair 25OHD synthesis via VDR mediated mechanism (Drincic *et al.*,2012). Despite the consensus of many previous observational studies that linked VDD to obesity (Goldner *et al.*, 2008), RCTs findings were contradictory about the effect of VD₃ on obesity in people with VDD or VD insufficiency (Turer *et al.*,2012). However, in the overwhelming majority of those trials have shown a positive effect of VD₃ on 25OHD and leptin. Independent association between 25OHD and leptin levels from adiposity was observed after a 1-year lifestyle intervention (Gangloff *et al.*, 2019). These observations may underscore the role of lifestyle modifications to decrease leptin levels during clinical management of VDD. Nevertheless, there is no certainty that the modulatory effect of VD₃ supplementation on leptin is involved in the improvement of insulin resistance in susceptible individuals to T2DM. Overall, Taking into account the U-

shaped curve. (Kojima G *et al.*,2017) and based on the dose dependent hypothesis, it has supposed that when VD intake is below, the serum leptin level will be low and vice versa VD supplementation may cause raising serum leptin (Tarcin *et al.*, 2010; Ghavamzadeh and Mahdavi, 2014) as shown in this review. In conclusion, we suggest further clinical trials wither to confirm or negation of existence of diabetogenic or anti-diabetogenic effects for VD₃ supplementations.

ACKNOWLEDGEMENT

The authors are grateful to the Applied Science Private University (ASU), for the full support.

Disclosure

The authors report no conflicts of interest in this work.

REFERENCES

1. Smith, A. G., & Singleton, J. R. (2013). Obesity and hyperlipidemia are risk factors for early diabetic neuropathy. *Journal of Diabetes and Its Complications*,27(5), 436-442. doi:10.1016/j.jdiacomp.2013.04.003
2. Al-Shaer, A. H., Abu-Samak, M. S., Hasoun, L. Z., Mohammad, B. A., & Bashedi, I. A. (2019). Assessing the effect of omega-3 fatty acid combined with vitamin D3 versus vitamin D3 alone on estradiol levels: A randomized, placebo-controlled trial in females with vitamin D deficiency. *Clinical Pharmacology: Advances and Applications*,Volume 11, 25-37. Doi:10.2147/cpaa.s182927
3. Qatatsheh, A., Tayyem, R., Al-Shami, I., Al-Holy, M. and Al-rethaia, A. (2015). Vitamin D deficiency among Jordanian university students and employees. *Nutrition & Food Science*, 45(1), pp.68-82.
4. Chadt, A., Scherneck, S., Joost, H., & Al-Hasani, H. (2018, January 23). Molecular links between Obesity and Diabetes: "Diabesity" (K. R. Feingold, B. Anawalt, A. Boyce, G. Chrousos, K. Dungan, A. Grossman, *et al.*, Eds.). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/25905279>
5. Vimaleswaran, K., Berry, D., Lu, C., Tikkanen, E., Pilz, S., Hiraki, L., . . . Hyppönen, E. (2015). Causal Relationship between Obesity and Vitamin D Status: Bi-Directional Mendelian Randomization Analysis of Multiple Cohorts. *Nutritional Biochemistry*,145-169. doi:10.1201/b18536-11
6. Hajimohammadi, M., Shab-Bidar, S., & Neyestani, T. R. (2016). Vitamin D and serum leptin: A systematic review and meta-analysis of observational studies and randomized controlled trials. *European Journal of Clinical Nutrition*,71(10), 1144-1153. Doi:10.1038/ejcn.2016.245
7. Ruiz-Ojeda FJ, Anguita-Ruiz A, Leis R, Aguilera CM. Genetic Factors and Molecular Mechanisms of Vitamin D and Obesity Relationship. *Ann Nutr Metab*. 2018;73(2):89-99. doi: 10.1159/000490669. Epub 2018 Jul 6. PMID: 29982250
8. Yadav A, Kataria MA, Saini V, Yadav A. Role of leptin and adiponectin in insulin resistance. *Clin Chim Acta*. 2013 Feb 18;417:80-4. doi: 10.1016/j.cca.2012.12.007. Epub 2012 Dec 22. PMID: 23266767.
9. Kim, M., Na, W., & Sohn, C. (2013). Correlation between vitamin D and cardiovascular disease predictors in overweight and obese Koreans. *Journal of Clinical Biochemistry and Nutrition*,167-171. Doi:10.3164/jcfn.12-81

Circulatory Leptin Levels as a Key Link in the Potential Association between Insulin Resistance and Vitamin D Deficiency: A Review Article

- Dinca, M., Serban, M., Sahebkar, A., Mikhailidis, D. P., Toth, P. P., Martin, S. S., . . . Banach, M. (2016). Does vitamin D supplementation alter plasma adipokines concentrations? A systematic review and meta-analysis of randomized controlled trials. *Pharmacological Research*, 107, 360-371. [Doi:10.1016/j.phrs.2016.03.035](https://doi.org/10.1016/j.phrs.2016.03.035)
- Hewison, M. (2012). An update on vitamin D and human immunity. *Clinical Endocrinology*, 76(3), pp.315-325.
- Newberry SJ, Chung M, Shekelle PG, Booth MS, Liu JL, Maher AR, Motala A, Cui M, Perry T, Shanman R, Balk EM. Vitamin D and Calcium: A Systematic Review of Health Outcomes (Update). *Evid Rep Technol Assess (Full Rep)*. 2014 Sep;(217):1-929. [doi: 10.23970/AHRQEPERTA217](https://doi.org/10.23970/AHRQEPERTA217). PMID: 30313003.
- Kovacs, C.S., 2014. Bone development and mineral homeostasis in the fetus and neonate: roles of the calciotropic and phosphotropic hormones. *Physiological Reviews*, 94(4), pp.1143-1218.
- Mundy, G.R. and Guise, T.A., 1999. Hormonal control of calcium homeostasis. *Clinical Chemistry*, 45(8), pp.1347-1352.
- Hosseini-nezhad, A. and Holick, M. (2013). Vitamin D for Health: A Global Perspective. *Mayo Clinic Proceedings*, 88(7), pp.720-755.
- Wimalawansa, S. (2016). Non-musculoskeletal benefits of vitamin D. *The Journal of Steroid Biochemistry and Molecular Biology* Retrieved from <https://www.journals.elsevier.com/the-journal-of-steroid-biochemistry-and-molecular-biology>
- Ran Zhang, & Declan P Naughton. (2010, December 08). Vitamin D in health and disease: Current perspectives. Retrieved from <https://nutritionj.biomedcentral.com/articles/10.1186/1475-2891-9-65>
- Chen, T., Chimeh, F., Lu, Z., Mathieu, J., Person, K., Zhang, A., Kohn, N., Martinello, S., Berkowitz, R. and Holick, M. (2007). Factors that influence the cutaneous synthesis and dietary sources of vitamin D. *Archives of Biochemistry and Biophysics*, 460(2), pp.213-217.
- Teske, K.A., Bogart, J.W., Sanchez, L.M., Olivia, B.Y., Preston, J.V., Cook, J.M., Silvaggi, N.R., Bikle, D.D. and Arnold, L.A., 2016. Synthesis and evaluation of vitamin D receptor-mediated activities of cholesterol and vitamin D metabolites. *European Journal of Medicinal Chemistry*, 109, pp.238-246.
- Nair, R. and Maseeh, A., 2012. Vitamin D: The "sunshine" vitamin. *Journal of Pharmacology and Pharmacotherapeutics*, 3(2), p.118.
- Jovičić, S., Ignjatović, S. and Majkić-Singh, N., 2012. Biochemistry and metabolism of vitamin D/Biohemija i metabolizam itamin D. *Journal of Medical Biochemistry*, 31(4), pp.309-315.
- Powe, C.E., Evans, M.K., Wenger, J., Zonderman, A.B., Berg, A.H., Nalls, M., Tamez, H., Zhang, D., Bhan, I., Karumanchi, S.A. and Powe, N.R., 2013. Vitamin D-binding protein and vitamin D status of black Americans and white Americans. *New England Journal of Medicine*, 369(21), pp.1991-2000.
- Lerchbaum, E. and Obermayer-Pietsch, B. (2012). MECHANISMS IN ENDOCRINOLOGY: Vitamin D and fertility: a systematic review. *European Journal of Endocrinology*, 166(5), pp.765-778.
- Theodoratou, E., Tzoulaki, I., Zgaga, L. and Ioannidis, J. (2014). Vitamin D and multiple health outcomes: umbrella review of systematic reviews and meta-analyses of observational studies and randomised trials. *BMJ*, 348(apr01 2), pp.g2035-g2035.
- Mahmoud Abu -Samak, Rula Khuzaie, Mohammad Abu-Hasheesh, Malak Jaradeh and Mohammad Fawzi, 2008. Relationship of Vitamin B12 Deficiency with Overweight in Male Jordanian Youth. *Journal of Applied Sciences*, 8: 3060-3063.
- Pearce, S.H. and Cheetham, T.D., 2010. Diagnosis and management of vitamin D deficiency. *Bmj*, 340(jan11 1), pp.b5664-b5664
- Gonzalez, L. (2014). Is vitamin D deficiency a major global public health problem? *The Journal of Steroid Biochemistry and Molecular Biology*, 144, pp.138-145.4, pp.138-145.
- Abu-Samak MS, AbuRuz ME, Masa'Deh R, Khuzai R, Jarrah S. Correlation of selected stress associated factors with vitamin D deficiency in Jordanian men and women. *Int J Gen Med*. 2019 Jun 28;12:225-233. [doi: 10.2147/IJGM.S198175](https://doi.org/10.2147/IJGM.S198175). PMID: 31303782; PMCID: PMC6612048.
- Mallah, E., Hamad, M., ElManaseer, M., Qinna, N., Idkaidek, N., Arafat, T., and Matalka, K. (2011). Plasma concentrations of 25-hydroxyvitamin D among Jordanians: Effect of biological and habitual factors on vitamin D status. *BMC Clinical Pathology*, 11(1).
- Gudjinu, H. Y., & Sarfo, B. (2017). Risk factors for type 2 diabetes mellitus among out-patients in Ho, the Volta regional capital of Ghana: A case-control study. *BMC Research Notes*, 10(1). [Doi:10.1186/s13104-017-2648-z](https://doi.org/10.1186/s13104-017-2648-z)
- Abu-Hasheesh MO, Abu-Samak MS, Al-Matubsi HY, et al. (2010). Association of parental history of type 2 diabetes mellitus with leptin levels in Jordanian male youth. *Saudi Med J*. 31: 882-6.
- Abu-Samak MS, Mohammad BA, Abu-Taha MI, Hasoun LZ, Awwad SH. Associations between sleep deprivation and salivary testosterone levels in male university students: a prospective cohort study. *Am J Mens Health*. 2018;12(2):411-419. [doi:10.1177/1557988317735412](https://doi.org/10.1177/1557988317735412)
- Abu-Taha M, Dagash R, Mohammad BA, Basheiti I, Abu-Samak MS. Combined Effect Of Coffee Consumption And Cigarette Smoking On Serum Levels Of Vitamin B12, Folic Acid, And Lipid Profile In Young Male: A Cross-Sectional Study. *Int J Gen Med*. 2019;12:421-432 <https://doi.org/10.2147/IJGM.S213737>
- Abu-Samak, M., Abu-Zaiton, A., Al-Jaberi, A., Sundookah, A., Atrooz, O., khadra, K., Kuzaie, R., & H. Talib, W. (2013). Morning Salivary Cortisol Associates with Elevated Serum Leptin Levels in Jordanian Young Men with Olive Pollen Induced Allergic Rhinitis. *Journal of Advances in Medicine and Medical Research*, 4(3), 797-806. <https://doi.org/10.9734/BJMMR/2014/5418>
- Gao, Z., Li, Q., Wu, X., Zhao, X., Zhao, L., & Tong, X. (2017). New Insights into the Mechanisms of Chinese Herbal Products on Diabetes: A Focus on the "Bacteria-Mucosal Immunity-Inflammation-Diabetes" Axis. *Journal of Immunology Research*, 2017, 1-13. [Doi:10.1155/2017/1813086](https://doi.org/10.1155/2017/1813086).

36. Jarab, A. S., Mukattash, T. L., Al-Azayzih, A., & Khdour, M. (2018). A focus group study of patient's perspective and experiences of type 2 diabetes and its management in Jordan. *Saudi Pharmaceutical Journal*,26(3), 301-305. [Doi:10.1016/j.jsps.2018.01.013](https://doi.org/10.1016/j.jsps.2018.01.013)
37. Tong, H. V., Luu, N. K., Son, H. A., Hoan, N. V., Hung, T. T., Velavan, T. P., & Toan, N. L. (2016). Adiponectin and pro-inflammatory cytokines are modulated in Vietnamese patients with type 2 diabetes mellitus. *Journal of Diabetes Investigation*,8(3), 295-305. [doi:10.1111/jdi.12579](https://doi.org/10.1111/jdi.12579)
38. González, F. (2012). Inflammation in Polycystic Ovary Syndrome: Underpinning of insulin resistance and ovarian dysfunction. *Steroids*,77(4), 300-305. [Doi:10.1016/j.steroids.2011.12.003](https://doi.org/10.1016/j.steroids.2011.12.003).
39. Ferguson, A. V. (2014). Circumventricular Organs: Integrators of Circulating Signals Controlling Hydration, Energy Balance, and Immune Function (L. A. De Luca, J. V. Menani, & A. K. Johnson, Eds.). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/24829996>
40. Lanzerstorfer, P., Yoneyama, Y., Hakuno, F., Müller, U., Höglinger, O., Takahashi, S., & Weghuber, J. (2015). Analysis of insulin receptor substrate signaling dynamics on microstructured surfaces. *The FEBS Journal*,282(6), 987-1005. [Doi:10.1111/febs.13213](https://doi.org/10.1111/febs.13213)
41. Finucane, F. M., Luan, J., Wareham, N. J., Sharp, S. J., O'Rahilly, S., Balkau, B., . . . Savage, D. B. (2009). Correlation of the leptin:adiponectin ratio with measures of insulin resistance in non-diabetic individuals. *Diabetologia*,52(11), 2345-2349. [Doi:10.1007/s00125-009-1508-3](https://doi.org/10.1007/s00125-009-1508-3).
42. Dsouza, A. M., Asadi, A., Johnson, J. D., Covey, S. D., & Kieffer, T. J. (2014). Leptin Deficiency in Rats Results in Hyperinsulinemia and Impaired Glucose Homeostasis. *Endocrinology*,155(4), 1268-1279. [Doi:10.1210/en.2013-1523](https://doi.org/10.1210/en.2013-1523)
43. Silva, A. A., Hall, J. E., & Carmo, J. M. (2017). Leptin reverses hyperglycemia and hyperphagia in insulin deficient diabetic rats by pituitary-independent central nervous system actions. *Plos One*,12(11). [doi:10.1371/journal.pone.0184805](https://doi.org/10.1371/journal.pone.0184805)
44. Frayn, K. N. (2001). Adipose tissue and the insulin resistance syndrome. *Proceedings of the Nutrition Society*,60(3), 375-380. [Doi:10.1079/pns200195](https://doi.org/10.1079/pns200195).
45. Klötting, N., & Blüher, M. (2014). Adipocyte dysfunction, inflammation and metabolic syndrome. *Reviews in Endocrine and Metabolic Disorders*,15(4), 277-287. [Doi:10.1007/s11154-014-9301-0](https://doi.org/10.1007/s11154-014-9301-0)
46. Khawaja, N., Liswi, M., El-Khateeb, M., Hyassat, D., Bajawi, D., Elmohtaseb, M., Alkhateeb, H. and Ajlouni, K. (2017). Vitamin D Dosing Strategies among Jordanians with Hypovitaminosis D. *Journal of Pharmacy Practice*, 30(2), pp.172-179.
47. Lagunova, Z., Porojnicu, A.C., Lindberg, F., Hexeberg, S. and Moan, J., (2009). The dependency of vitamin D status on body mass index, gender, age and season. *Anticancer Research*, 29(9), pp.3713-3720.
48. Konradsen, S., Ag, H., Lindberg, F., Hexeberg, S. and Jorde, R. (2008). Serum 1,25-dihydroxy vitamin D is inversely associated with body mass index. *European Journal of Nutrition*, 47(2), pp.87-91.
49. Mead, M. N. (2008). Benefits of Sunlight: A Bright Spot for Human Health . *Environmental Health Perspectives*,116(4). [Doi:10.1289/ehp.116-a160](https://doi.org/10.1289/ehp.116-a160)
50. Pittas, A. G., Dawson-Hughes, B., Sheehan, P., Ware, J. H., Knowler, W. C., Aroda, V. R., . . . Staten, M. (2019). Vitamin D Supplementation and Prevention of Type 2 Diabetes. *New England Journal of Medicine*. [Doi:10.1056/nejmoa1900906](https://doi.org/10.1056/nejmoa1900906)
51. Boucher, B. J., John, W. G., & Noonan, K. (2004). Hypovitaminosis D is associated with insulin resistance and β cell dysfunction. *The American Journal of Clinical Nutrition*,80(6), 1666-1666. [Doi:10.1093/ajcn/80.6.1666](https://doi.org/10.1093/ajcn/80.6.1666)
52. Gandhe, M. B. (2013). Evaluation of 25(OH) Vitamin D 3 with Reference to Magnesium Status and Insulin Resistance in T2DM. *Journal Of Clinical And Diagnostic Research*. [Doi:10.7860/jcdr/2013/6578.3568](https://doi.org/10.7860/jcdr/2013/6578.3568)
53. Upreti, V., Maitri, V., Dhull, P., Handa, A., Prakash, M., & Behl, A. (2018). Effect of oral vitamin D supplementation on glycemic control in patients with type 2 diabetes mellitus with coexisting hypovitaminosis D: A parallel group placebo controlled randomized controlled pilot study. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*,12(4), 509-512. [doi:10.1016/j.dsx.2018.03.008](https://doi.org/10.1016/j.dsx.2018.03.008)
54. Monnier, L., & Colette, C. (2010). Vitamin D and diabetes: Much ado about nothing? *Diabetes & Metabolism*,36(5), 323-325. [Doi:10.1016/j.diabet.2010.06.003](https://doi.org/10.1016/j.diabet.2010.06.003)
55. Niroomand, M., Fotouhi, A., Irannejad, N., & Hosseiniapanah, F. (2019). Does high-dose vitamin D supplementation impact insulin resistance and risk of development of diabetes in patients with pre-diabetes? A double-blind randomized clinical trial. *Diabetes Research and Clinical Practice*,148, 1-9. [Doi:10.1016/j.diabres.2018.12.008](https://doi.org/10.1016/j.diabres.2018.12.008)
56. Knekt, P., Laaksonen, M., Mattila, C., Härkänen, T., Marniemi, J., Heliövaara, M., . . . Reunanen, A. (2008). Serum Vitamin D and Subsequent Occurrence of Type 2 Diabetes. *Epidemiology*,19(5), 666-671. [Doi:10.1097/ede.0b013e318176b8ad](https://doi.org/10.1097/ede.0b013e318176b8ad)
57. Forsythe, L. K., Livingstone, M. B., Barnes, M. S., Horigan, G., Mccorley, E. M., Bonham, M. P., . . . Wallace, J. M. (2011). Effect of adiposity on vitamin D status and the 25-hydroxycholecalciferol response to supplementation in healthy young and older Irish adults. *British Journal of Nutrition*,107(1), 126-134. [Doi:10.1017/s0007114511002662](https://doi.org/10.1017/s0007114511002662).
58. Harroud, A., & Richards, J. B. (2018). Mendelian randomization in multiple sclerosis: A causal role for vitamin D and obesity? *Multiple Sclerosis Journal*,24(1), 80-85. [Doi:10.1177/1352458517737373](https://doi.org/10.1177/1352458517737373)
59. Lenders, C. M., Feldman, H. A., Scheven, E. V., Merewood, A., Sweeney, C., Wilson, D. M., . . . Holick, M. F. (2009). Relation of body fat indexes to vitamin D status and deficiency among obese adolescents. *The American Journal of Clinical Nutrition*,90(3), 459-467. [Doi:10.3945/ajcn.2008.27275](https://doi.org/10.3945/ajcn.2008.27275)
60. Duggan, C., Tapsoba, J. D., Mason, C., Imayama, I., Korde, L., Wang, C., & Mctiernan, A. (2015). Effect of Vitamin D3 Supplementation in Combination with Weight Loss on Inflammatory Biomarkers in Postmenopausal Women: A Randomized Controlled Trial. *Cancer Prevention Research*,8(7), 628-635. [Doi:10.1158/1940-6207.capr-14-0449](https://doi.org/10.1158/1940-6207.capr-14-0449)

Circulatory Leptin Levels as a Key Link in the Potential Association between Insulin Resistance and Vitamin D Deficiency: A Review Article

61. Mousa, A., Naderpoor, N., Wilson, K., Plebanski, M., Courten, M. P., Scragg, R., & Courten, B. D. (2019). Vitamin D supplementation increases adipokine concentrations in overweight or obese adults. *European Journal of Nutrition*. Doi:10.1007/s00394-019-01899-5
62. Vahdat, S., Naini, A., Hedaiati, Z., Shahzeidi, S., Pezeshki, A., & Nasri, H. (2016). The effect of vitamin D administration on serum leptin and adiponectin levels in end-stage renal disease patients on hemodialysis with vitamin D deficiency: A placebo-controlled double-blind clinical trial. *Journal of Research in Medical Sciences*,21(1), 1. doi:10.4103/1735-1995.175144
63. Mai, S., Walker, G., Vietti, R., Cattaldo, S., Mele, C., Priano, L., . . . Marzullo, P. (2017). Acute Vitamin D3 Supplementation in Severe Obesity: Evaluation of Multimeric Adiponectin. *Nutrients*,9(5), 459. Doi:10.3390/nu9050459.
64. Entezari, M., Khosravi, Z., Kafeshani, M., Tavasoli, P., & Zadeh, A. (2018). Effect of Vitamin D supplementation on weight loss, glycemic indices, and lipid profile in obese and overweight women: A clinical trial study. *International Journal of Preventive Medicine*,9(1), 63. Doi:10.4103/ijpvm.ijpvm_329_15
65. Mott, J. W., Wang, J., Thornton, J. C., Allison, D. B., Heymsfield, S. B., & Pierson, R. N. (1999). Relation between body fat and age in 4 ethnic groups. *The American Journal of Clinical Nutrition*,69(5), 1007-1013. Doi:10.1093/ajcn/69.5.1007
66. Abu-Samak, M., Yousef, A., Al-Jarie, A., Al-Matubsi, H. Y., Abu-Zaiton, A., Al-Quraan, M., & Khuzaie, R. (2011). Lipid and Hematological Parameters in Hyperleptinemic Healthy Arab Male Youth in Jordan. *Pakistan Journal of Biological Sciences*,14(5), 344-350. Doi:10.3923/pjbs.2011.344.350
67. Wortsman, J., Matsuoka, L. Y., Chen, T. C., Lu, Z., & Holick, M. F. (2000). Decreased bioavailability of vitamin D in obesity. *The American Journal of Clinical Nutrition*,72(3), 690-693. doi:10.1093/ajcn/72.3.690
68. Targher G, Bertolini L, Scala L, Cigolini M, Zenari L, Falezza G, et al. (2007) Associations between serum 25-hydroxyvitamin D3 concentrations and liver histology in patients with non-alcoholic fatty liver disease. *Nutr Metab Cardiovasc Dis*. 17:517-24. doi: 10.1016/j.numecd.2006.04.002
69. Drincic AT, Armas LA, Van Diest EE, Heaney RP. (2012) Volumetric dilution, rather than sequestration best explains the low vitamin D status of obesity. *Obesity* 20:1444-8. Doi: 10.1038/oby.2011.404
70. Goldner, W. S., Stoner, J. A., Thompson, J., Taylor, K., Larson, L., Erickson, J., & McBride, C. (2008). Prevalence of Vitamin D Insufficiency and Deficiency in Morbidly Obese Patients: A Comparison with Non-Obese Controls. *Obesity Surgery*,18(2), 145-150. Doi:10.1007/s11695-007-9315-8
71. Turer, C. B., Lin, H., & Flores, G. (2012). Prevalence of Vitamin D Deficiency Among Overweight and Obese US Children. *Pediatrics*,131(1). doi:10.1542/peds.2012-1711
72. Gangloff, A., Bergeron, J., Lemieux, I., Tremblay, A., Poirier, P., Alméras, N., & Després, J. (2019). Relationships between circulating 25(OH) vitamin D, leptin levels and visceral adipose tissue volume: Results from a 1-year lifestyle intervention program in men with visceral obesity. *International Journal of Obesity*. Doi:10.1038/s41366-019-0347-7.
73. Kojima G, Iliffe S, Tanabe M. Vitamin D supplementation as a potential cause of U-shaped associations between vitamin D levels and negative health outcomes: a decision tree analysis for risk of frailty. *BMC Geriatr*. 2017 Oct 16;17(1):236. doi: 10.1186/s12877-017-0631-0. PMID: 29037174; PMCID: PMC5644251.
74. Tarcin, O., Yavuz, D. G., Ozben, B., Telli, A., Ogunc, A. V., Yuksel, M., . . . Akalin, S. (2010). Effect of Vitamin D Deficiency and Replacement on Endothelial Function in Asymptomatic Subjects. *Obstetrical & Gynecological Survey*,65(3), 173-174. doi:10.1097/01.ogx.0000369678.64828.55
75. Saeid Ghavamzadeh, M. M., and Reza Mahdavi. (2014 Sep 5). The Effect of Vitamin D Supplementation on Adiposity, Blood Glycated Hemoglobin, Serum Leptin and Tumor Necrosis Factor- α in Type 2 Diabetic Patients. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4192769/>