

EFFECT OF GIVING ETHANOL JENKOL SEEDS (*Pithecellobium lobatum* Benth.) ON THE IMPROVEMENT OF UREUM AND CREATININE LEVELS OF WHITE RAT

Evi Kurniawaty^a, Frisca Febe Lumban Gaol^b, Intanri Kurniati^c, Silvia Andriani^d, Soraya Rahmanisa^e, Silvia Andriani^f

^{a,b,c,d,e,f}University of Lampung, Lampung Indonesia

Corresponding author: evikurniawatyherlambang@yahoo.com 1 *, co-authors: silviaandriani12@gmail.com (evikurniawaty@fk.unila.ac.id)

Abstract

Purpose: Diabetes Mellitus is a degenerative disease which is a chronic disease caused by the failure of the pancreas to produce insulin. Diabetes Mellitus is characterized by polyuria, polydipsia, and polyphagia, accompanied by an increase in blood glucose. Diabetes mellitus triggers many complications, one of which is kidney problems. Jengkol is a type of plant that can lower blood sugar levels, but it can also cause acute kidney failure due to the jengkollic acid content in it. To diagnose acute renal failure, it is necessary to examine kidney function by measuring levels of urea and creatinine.

Research Methodology: This research is an experimental study with a Post Test Only Control Group Design. The sample in this study were male white rats (*Rattus norvegicus*) Sprague Dawley strain with a body weight of 200-250 grams, aged 3-4 months, totaling 25 animals, which were divided into 5 groups.

Results: Based on the results of the oneway ANOVA test, it was found that the p value > 0.05 jengkol seed extract had no effect on the increase in urea and creatinine levels of sprague dawley rats induced by alloxan. However, when viewed from the mean creatine level at a dose of 1200 mg / kgbb, the ethanol extract of jengkol seeds can cause an increase in creatinine levels with a group mean value of 0.08mg / dl.

Limitations: This research is expected to be the basis for further researchers regarding jengkol extract as a therapy for kidney failure

Contribution: Bandar lampung, Biologi Molekuler

Keywords: Alloxan, Glucose, creatinine, and polyura.

1. INTRODUCTION

Diabetes Mellitus is one of the non-communicable diseases that will increase in number in the future. This is presumably because the people's lifestyle in consuming a lot of foods that contain protein, fat, sugar, salt and a little fiber. Suyono, 2010. One type of plant that can be used as a diabetes mellitus drug is the jengkol plant. The jengkol plant contains several chemical compounds such as flavonoids, tannins and saponins. Flavonoids have antioxidants.

2. LITERATURE REVIEW AND HYPOTHESIS DEVELOPMENT

According to research (Elysa, 2011) jengkol seeds have an effect on reducing glucose levels in mice, but it does not explain whether the dose used is safe and does not cause toxicity. Jengkol poisoning is acute kidney failure. Jengkolat acid contained in jengkol can form crystals that settle and cause obstruction in Jha et al. 2008. Obstruction in the urinary tract is one of the causes of acute kidney failure. To confirm the diagnosis of acute renal failure, it is necessary to examine renal function, namely examining the levels of serum urea and creatinine Molitoris 2009.

In diabetics the incidence rate for cases of acute complications is still very high. Acute complications in diabetes are diabetic ketoacidosis (DKA) and hyperglycaemic hyperosmolarity (HHS). Chronic complications from diabetes can include

vascular complications, which are divided into microvascular and macrovascular. Macrovascular, namely coronary artery disease, lower limb blood vessels and macrovascular, namely retinopathy, nephropathy, and others. (Waspadji, 2010).

3. RESEARCH METHODOLOGY

This research is an experimental research with the design of Post Test Only Control Group Design. A sample of 25 adult male white rats (*Rattus norvegicus*) Sprague Dawley strain for each treatment, aged 3- 4 months, weighing 200 - 2500 grams. Divided into 5 treatment groups, namely 1). In the negative control group, rats were only given distilled water orally without alloxan induction. In the second group the mice were given distilled water orally and alloxan induced intraperitoneally. The third group was treated with the ethanol extract of jengkol seeds with a dose of 600 mg / kgbb orally and induced by alloxan. In the fourth group treated with the ethanol extract of jengkol seeds at a dose of 900 mg / kgbb orally and induced by alloxan. In the fifth group treated with the ethanol extract of jengkol seeds at a dose of 1200 mg / kgbb orally and induced by alloxan. The treatment was given for 14 days. Blood samples were taken at the end of the study. The blood sample was collected using a red top vacuotainer tube, centrifuged for 10 minutes at a speed of 4000 rpm, then the urea and serum creatinine were examined by spectrophotometry.

Effect Of Giving Ethanol Jengkol Seeds (*Pithecellobium Lobatum Benth.*) On The Improvement Of Ureum And Creatinine Levels Of White Rat

of Health's creatinine levels.

4. RESULTS AND DISCUSSIONS

Table 1. One-way ANOVA test results blood glucose levels

	Sum of	Df	Mean	F	Sig
	Squares		Square		
Between Groups	1204,400	4	301,100	2491	,076
Within Groups	2417,600	20	120,880		
Total	3622,000	24			

In the ANOVA test results table, the p value is obtained = 0.076, which means there is no difference in urea levels between groups.

Table 2. Urea Levels Normality Test Results

GROUP	Shapiro-Wilk			
	Statistics	df	Sig.	
Ureum	I	,927	5	,573
	II	,834	5	,149
	III	,816	5	,109
	IV	,928	5	,586
	V	,885	5	,332

In the table of normality test results, the p value for the five groups is > 0.05, so it can be concluded that the distribution of the three groups is normal.

Table 3. One-way ANOVA test results creatinine levels

	Sum of	Df	Mean	F	Sig
	Squares		Square		
Between Groups	1204,400	4	301,100	2491	,076
Within Groups	2417,600	20	120,880		
Total	3622,000	24			

Table 4. Result of Normality Test of Creatinine Levels

GROUP	Shapiro-Wilk			
	Statistics	df	Sig.	
Creatinine	I	,987	5	,967
	II	,961	5	,814
	III	,932	5	,607
	IV	,956	5	,777
	V	,961	5	,814

In the table of normality test results, the p value for the five groups is > 0.05, so it can be concluded that the distribution of the three groups is normal.

The results showed that the treatment group was not so significant, with a p value > 0.05, meaning that there was no significant difference in urea levels between the study groups. Ureum is not an ideal index of renal function because many events outside the kidney affect plasma levels. However, examination of plasma urea levels is still important and necessary in patients with kidney disease, especially to evaluate the effect of dietary protein restriction.

The results of creatinine levels in the treatment group and 2 control groups did not show any significant difference. It can be seen that the value of p > 0.05 means that there is no difference in creatinine levels between groups. However, if we look at the creatinine level in group 5, it has passed the limit of normal creatinine levels in mice with a value of 0.86. This indicates that a dose of 1200 mg / kgbb causes a decrease in renal function as indicated by an increase in creatinine levels > 0.3 mg / dl.

Several factors that can affect creatinine levels are gender, hunger conditions, and Guyton and Hall 2007 muscle tissue size. In addition, strenuous exercise and surgical procedures that damage skeletal muscles can increase the 2011 Ministry

5. CONCLUSION

Giving jengkol seed ethanol extract at a dose of 1200 mg / kgbb can cause a decrease in renal function, which is indicated by an increase in urea levels > 0.3 mg / dl in Sparague Dawley rats induced by alloxan

LIMITATION AND STUDY FORWARD

This research is expected to be the basis for further researchers regarding jengkol extract as a therapy for kidney failure

ACKNOWLEDGEMENT

The author would like to thank the Faculty of Medicine, University of Lampung for fully funding this research through DIPA FK UNILA 2020.

REFERENCES

1. Adewani N. 2008. Pengaruh pemberian Rebusan Kulit Kayu Duwet Terhadap Presentase Penurunan Kadar Glukosa Darah pada Tikus Putih. *Skripsi*. Yogyakarta: Universitas Muhammadiyah Yogyakarta.
2. Astrian RT. 2009. Pengaruh Antioksidan Polifenol terhadap Kadar Gukosa Darah dan Insulin Mencit (*Mus musculus L.*) S.W. jantan yang dikondisikan Diabetes Mellitus. *Skripsi*. Jawa Barat. Institut teknologi Bandung.
3. Atmajaya Y. 2012. Perbandinagn Efek Hipoglikemik Biji Petai Cina (*Lucaena luchocephala*) dengan Glibenklamik pada tikus putih yang diinduksi dengan Alloxan. *Journal Mutiara Medika* 2012 8(9)
4. Busatta F. 2011. Obesity, Diabetes an the Thrifty Gene. *Antrocom Online Journal of Anthropology* 2011 7 (1).
5. Dahalan MS. 2011. *Statistik Untuk Kedokteran dan Kesehatan Edisi Kelima*. Jakarta: Salemba Medika.
6. Elysa. 2011. Uji Efek Ekstrak Etanol Biji Jengkol (*Pithecellobium lobatum Benth.*) terhadap penurunan kadar glukosa darah tikus putih jantan galur Wistar yang diinduksi aloksan. (*Skripsi*). Medan: Fakultas Farmasi Universitas Sumatera Utara
7. Mikito A, Yamashita C, Iwasaki Y, 1995. A Triterpenoid Saponin Ectraxtion, there of and use Treat or Prevent Diabetes Meliitus. *European Patient Application*.
8. Muthusamy VS, Anand S, Sangeetha KN, Sujatha S, Balakrishnan A, Lakshmi BS. 2008. Tannis Presents in Chicorium Intybus enchaneGlucose uptake and inhibit adipogenesis in 3T3-L1 adipocytes trough PTB1B inhibition. *Chemico –Biological Interaction* 174 (1)
9. Ramadhani R. 2008. Pengaruh Ekstrak Etanol Daun Muntingia calabura L. terhadap Kadar Glukosa Darah Mencit (*Mus musculus L.*) Swiss Webster Jantan Dewasa yang dikondisikan. *Skripsi*. Jawa Barat: Institut Teknologi Bandung.
10. Song Y, JoAnn EM, Julie EB, Howard DS, Simin L, 2005. Associations of Dietary Flavonnoids with Riskof Type 2 Diabetes, and Markers of Insulin Resistance and Systemic Inflammation in WomenA Prospective Study and Cross Sectional Analysis. *Journal of the American College of nutrition* 24(5).
11. Sudoyo AW, Setyohadi B, Alwi I, Simadibrata KM, Setiati S. 2010. *Buku Ajar Ilmu Penyakit Dalam Edisi Kelima jilid 3*. Jakarta: Balai Penerbit Fakultas Kedokteran Universitas Indonesia.
12. Waspadji S. 2010. *Buku Ajar Ilmu Penyakit Dalam Edisi Kelima Jilid Ketiga*. Jakarta: Balai Penerbit Fakultas Kedokteran Universitas Indonesia.