

Effect of Protecting Proteins from Degradation in the Rumen on Rumen Fermentations of Al Awassi Lambs

Dr. Ibrahim S. Jasim

Prime Minister Advisory Commission, Baghdad, Iraq.

Email: newtrench259@yahoo.com

ABSTRACT

The present experiment was carried out to investigate the effect of different percentages (50 and 100%) of dried whey powder and sun flower meal treated with blood or formaldehyde on some Rumen Fermentations: Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃) and the total count of anaerobic bacteria (BC), in lambs fattening diets. The results showed significant decrease ($P < 0.05$) in (pH) and insignificant effect in (TVFA), (NH₃) and (BC) for dried whey powder treated with blood or formaldehyde compared sun flower meal treated with blood or formaldehyde, While There was insignificant effect in (pH), (TVFA), (NH₃) and (BC) for dried whey powder treated with blood in percentages 100% compared sun flower meal treated with blood in percentages 100%, and insignificant effect in (pH), (TVFA), (NH₃) and (BC) for dried whey powder treated with blood in percentages 50% compared sun flower meal treated with blood in percentages 50%, and insignificant effect in (pH), (TVFA), (NH₃) and (BC) for dried whey powder treated with formaldehyde in percentages 100% compared sun flower meal treated with formaldehyde in percentages 100%, and insignificant effect in (pH), (TVFA), (NH₃) and (BC) for dried whey powder treated with formaldehyde in percentages 50% compared sun flower meal treated with formaldehyde in percentages 50%.

Keywords: dried whey powder, sunflower meal, blood, formaldehyde, Rumen Fermentations.

Correspondence:

Dr. Ibrahim S. Jasim

Prime Minister Advisory Commission, Baghdad, Iraq.

Email: newtrench259@yahoo.com

INTRODUCTION

The degradable protein in the rumen supplies the rumen microorganisms with NH₃-N, amino acids and peptides. The decrease in the level of degradable protein in the rumen will lead to a decrease in the performance of the animal, as there was an increase in the efficiency of the nutritional conversion and the daily weight increase for the increase in the degradable protein in the rumen (Cooper *et al.*, 2002). Emanuele and Putnam (2006) show that the rapid decrease in the level of NH₃-N in the rumen indicates either the rapid growth of rumen microorganisms and their consumption of NH₃-N or their absorption through the rumen wall, that the concentration of NH₃-N and total volatile fatty acids TVFA increases linearly with increasing protein. The degradation in the rumen (Dhali *et al.*, 2006). Also, increasing the hydrolyzed protein in the rumen beyond the permissible limit in the ruminant diets increases the level of NH₃-N in the rumen, which leads to an increase in the blood (Nisa *et al.*, 2008). The rumen microbial ecosystem consists mainly of species of bacteria, fungi and protozoa, in quantities ranging from 10⁹ to 10¹¹ cfu / mL, and the diversity of these species and their proportions within the microbial community depend primarily on the diet consumed by ruminants (Fébel & Fekete, 1996). Studies report that rumen microbial communities have a high ability to degrade protein (Ørskov, 1992; Bach *et al.*, 2005; Nolan & Dobos, 2005), as *Bacteroides rumenicola* and *Peptostreptococcus* sp. It has a high ability to decompose protein in the rumen, synergistic action of different types of proteases producing peptides and amino acids (Wallace *et al.*, 1997). The pH is an important factor in the degradation of protein sources in the rumen, and its decrease leads to a decrease in the level of growth in the rumen microorganism and this negatively affects the fermentation and digestion of cellulose and the level of protein degradation decreases with the decrease in the

pH level in the rumen environment (Strobel & Russel, 1986; Cardozo *et al.*, 2000; 2002).

The increased need for animal production prompted researchers to use oilseed grains to feed ruminants as protein sources such as *Helianthus annuus*, an important oil crop characterized by high protein content with high sulfuric acid content (Daghir *et al.*, 1980). The third largest source of protein used for ruminants feed after soybean and canola seedling (USDA-FAS, 2017). The protein of the sunflower is characterized by its solubility and high decomposition compared to the other protein sources. Therefore, there are obstacles to meet the needs of highyielding dairy cows, calves and fast-growing sheep because the protein is rapid decomposition in the rumen, producing peptides, amino acids and ammonia, which reduces the degree of utilization and loss of amino acids and low digestibility (Lusus, 1982). Whey was considered a non-conventional, fastdegradable protein source, it is a byproduct of cheese making process of milk, containing 7% solid materials consisting of 4.9% lactose, 0.6% ash, low amounts of fat acid and protein (15-20%) and most whey is eliminated as a neglected product, so the challenge for nutritionists is to find the best way to benefit from it (El-shewy, 2016). In the low-protein feed, substituting the urea substitutes for improved urea performance compared with the soybean meal with urea, which resulted in less improvement in animal performance. The addition of shark also increased the production of microbial protein and improved feed utilization (Stock *et al.*, 1986). Research in livestock feed in many countries has shown that straw as a byproduct of cheese production can be used to feed large ruminants without any negative effects. There are also studies on determining optimal levels of addition, taking into account the benefits that will be achieved by limiting use of concentrates and disposal as an accidental product for dairy manufacturers and environmental pollution prevention (Salem *et al.*, 2007).

OBJECTIVE OF THE STUDY

Study of the effect of replacing the dried whey powder treated with blood or formaldehyde, sun flower meal treated with blood or formaldehyde and effect of replacing different percentages (50 and 100%) of dried whey powder and sun flower meal treated with blood or formaldehyde in the fattening diets on some Rumen Fermentations : Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N) and The total count of anaerobic bacteria (BC), of Al Awassi Lambs.

MATERIALS AND METHODS

This study was conducted in the animal field of the Animal Production Department, Faculty of Agricultural Engineering Sciences, University of Baghdad. The experiment lasted for 60 days preceded by a preliminary period of 14 days for the period from 2 of December 2017 to 13 of February 2018.

Preparation of feed materials

All raw materials, such as barley, wheat bran, dried whey powder, sunflower and dried whey powder, were purchased from the local markets. Random samples were taken for the purpose of conducting chemical analyzes and using the green alfalfa from the fields of the Faculty of Agricultural Engineering Sciences, University of Baghdad and conducting chemical analyzes (Table 1).

Treatment of the sunflower meal with fresh blood

Blood was collected from ruminants that were slaughtered in the Karkh massacre in containers containing citrate of sodium (6.8 g/L blood). The blood was then added to the sun flower by using an equal weight of blood and weight (1: 1) and then mixed by hand and dried in a fan oven at 60°C for 24 hours, after that, the sun flower was manually broken and packed in bags until it was used (Matsumoto *et al*, 1995).

Treatment of sunflower meal with formaldehyde

The sun flower was treated with 5% formaldehyde solution and 1 liter solution/10 kg dry matter from the sun flower by sprinkler after brushing the sun flower over a piece of nylon on the ground in a closed chamber with constant flipping to ensure that the solution reaches all parts of the sunflower to obtain a homogeneous level of treatment. The formaldehyde sunflower was kept in tightly sealed nylon bags and left for 72 hours for interaction between formaldehyde and sunflower meal. The bags and their contents were then emptied onto a nylon piece inside a well-ventilated hall for 48 hours to allow for the volatilization of the unformed formaldehyde solution, then the sunflower was put in bags until it was used (Hassan *et al*, 1990).

Treatment of dried whey powder with fresh blood

Blood was collected from the ruminants that were slaughtered in the Karkh massacre in containers containing citrate of sodium (6.8 g/L blood). The blood was then added to the dried whey powder using an equal weight of blood and dried whey powder by 1: 1 and then mix it byhand and dry it in a fan oven at 60°C for 24 hours. Then it was manually broken and packed in bags for use. (Matsumoto *et al*, 1995).

Treatment of dried whey powder with formaldehyde

Dried whey powder was treated with 5% formaldehyde solution and 1 liter solution/10 kg dry matter of dried whey powder by sprinkler after brushing the whey powder over a piece of nylon on the ground in a closed chamber with continuous stirring to ensure that the solution reached all parts of the whey powder to obtain a homogeneous level of treatment. The dried whey powder was stored in sealed nylon bags were left for 72 hours for interaction between formaldehyde and whey powder. The bags and their contents were then emptied onto a nylon piece inside a well-ventilated hall for 48 hours to allow the volatilization of the Non-reacting formaldehyde and then dried whey powder was packed in bags until it was used (Hassan *et al*, 1990).

Table 1: Chemical composition of raw materials in the installation of concentrates and fresh grit based on dry matter (%).

Feeding materials Chemical composition %	Barley	Wheat barn	Sunflower treated with blood	Sunflower treated with formaldehyde	Whey treated with blood	Whey treated with formaldehyde	Fresh alfalfa
Dry matter	90.12	89.87	94.77	93.30	97.59	95.86	27.22
Organic matter	93.58	91.59	89.31	85.78	96.13	94.68	91.13
Crude protein	12.22	14.72	21.37	21.67	21.08	19.11	18.21
Crude fiber	5.72	10.11	15.35	15.55	---	---	27.15
Ether Extract	3.15	4.63	9.79	10.05	7.39	8.17	3.03
Ash	6.42	8.41	8.00	8.04	6.39	6.02	8.87
Nitrogen free extract	72.49	62.13	42.42	42.44	64.32	65.82	42.74
Acid detergent fiber	27.13	48.45	38.44	37.88	---	---	45.75
Neutral detergent fiber	6.27	14.24	26.92	27.50	---	---	33.91
Lignin	1.35	2.88	9.88	10.50	---	---	8.77
Cellulose	4.92	11.36	17.04	17.00	---	---	25.14
Hemicellulose	20.86	34.21	11.52	10.38	---	---	11.84
Metabolic energy (Mica Gul/kg)	12.7	12.3	12.7	12.7	14.1	14.2	10.2

Metabolic energy (Mg / kg of material as is) = 0.012 × crude protein + 0.031 x ether extract + 0.005 × raw fiber + 0.014 × nitrogen-free extract (Maff, 1975).

GROWTH EXPERIMENT

Animals and experiment design

Two experiments were done use 16 lambs (Al Awassi strain) were purchased from the local markets. The average age of the lambs was 5-6 months and the average weight was 23.87± 0.56 kg. The lambs were randomly divided into 4 treatments and 4 lambs per treatment. The experimental treatments involved treatment T1 and T2 treated with blood (dried whey powder , sun flower meal) with substitution ratios 50 and 100% while T3 and T4 treated with formaldehyde blood (dried whey powder , sun flower meal) with substitution ratios 50 and 100% To compare the significant differences between the averages with a test (T). The lambs were distributed in single pens with an area of 2 × 2m² for each treatment and numbered according to their own treatment.

Experimental diets

The animals were fed on the experimental diets and according to the treatments shown in Table (2,3). The dried whey powder treated with Whole blood 50% instead of the untreated sunflower in diet of (T1) and all the other components of the diet remain constant, The dried whey powder treated with Whole blood 100% instead of the untreated sunflower in diet of (T2) and all the other components of the diet remain constant, in diet of (T3) the dried whey powder treated with formaldehyde 50% instead of the untreated sunflower and all the other components of the diet remain constant, in diet of (T4) the dried whey powder treated with formaldehyde 100% instead of the untreated sunflower and all the other components of the diet remain constant, In the first experiment. In the second experiment The sunflower treated with Whole blood 50% instead of the untreated sunflower in diet of (T1) and all the other components of the diet remain constant, The sunflower treated with Whole blood 100% instead of the untreated sunflower in diet of (T2) and all the other components of the diet remain constant, in diet of (T3) the sunflower treated with formaldehyde 50% instead of the untreated sunflower and all the other components of the diet remain constant, in diet of (T4) the sunflower treated with formaldehyde 100% instead of the untreated sunflower and all the other components of the diet remain constant. The lambs were fed gradually for 14 days before the start of the experiment, the concentrated diet was served once daily at 8:00 am and by 3% of the body weight in addition the alfalfa was provided freely and separated from the concentrated feed while the amounts of concentrated feed based on the new body weight for each lamb were adjusted weekly. The lambs were weighed at the beginning of the experiment in a In a special scale to determine the primary weight and then the process of weighing on a weekly basis and before the morning ration to calculate the rate of daily weight increase and then the weight of lambs at the end of the experiment to determine the final weight, the remaining feed was collected from concentrated diet and alfalfa every morning and before morning ration to calculate the daily feed intake as well, clean water was provided continuously in special metal containers that are cleaned daily, the lambs were vaccinated against the internal and external parasites as the animals were vaccinated against hepatic worms and bariatric with the continued control of the confidentiality throughout the duration of the experiment.

Table 2: Percentage of the primary components involved in the composition of concentrates of first experiment (%).

Type of treatment	Treatment with blood		Treatment with formaldehyde	
Replacement ratio %	50	100	50	100
Treatments	T1	T2	T3	T4
Feeding materials				
Barley	45	45	45	45
Wheat bran	40	40	40	40
Sunflower meal	6.5	0	6.5	0
dried whey powder treated with blood	6.5	13	0	0
dried whey powder treated with formaldehyde	0	0	6.5	13
*Mix minerals and vitamins	2	2	2	2

*Mix minerals and vitamins table 6

Table 3: Percentage of the primary components involved in the composition of concentrates of second experiment (%).

Type of treatment	Treatment with blood		Treatment with formaldehyde	
Replacement ratio %	50	100	50	100
Treatments	T1	T2	T3	T4
Feeding materials				
Barley	45	45	45	45
Wheat bran	40	40	40	40
Sunflower meal	6.5	0	6.5	0
Sunflower treated with blood	6.5	13	0	0
Sunflower treated with	0	0	6.5	13

formaldehyde				
*Mix minerals and vitamins	2	2	2	2

*Mix minerals and vitamins table 6

Table 4: Chemical analysis of experimental treatments for first experiment based on dry matter.

Type of treatment	Treatment with blood		Treatment with formaldehyde	
Replacement ratio %	50	100	50	100
Treatments	T1	T2	T3	T4
Chemical composition				
Dry matter	98.41	98.45	98.14	97.03
Organic matter	92.74	93.52	93.85	93.66
Crude protein	14.53	15.04	15.28	15.44
Crude fiber	8.53	7.44	8.29	8.81
Ether Extract	5.09	4.52	5.10	5.56
Ash	7.26	6.48	6.14	6.34
Nitrogen free extract	64.69	66.52	65.18	63.85
Acid detergent fiber	36.01	35.25	35.20	35.61
Neutral detergent fiber	13.50	12.66	13.29	12.81
Lignin	2.41	2.12	2.50	2.18
Cellulose	11.09	10.54	10.79	10.63
Hemicellulose	22.51	22.59	21.91	22.80
Metabolic energy (Mica Gul/kg)	12.7	12.9	12.9	12.8

Metabolic energy (Mg / kg of material as is) = 0.012 × crude protein + 0.031 x ether extract + 0.005 × raw fiber + 0.014 × nitrogen-free extract (Maff, 1975).

Table 5: Chemical analysis of experimental treatments for Second experiment based on dry matter.

Type of treatment	Treatment with blood		Treatment with formaldehyde	
Replacement ratio %	50	100	50	100
Treatments	T1	T2	T3	T4
Chemical composition				
Dry matter	97.33	98.18	96.30	95.21
Organic matter	91.58	92.88	92.09	93.58
Crude protein	15.36	15.18	15.50	15.63
Crude fiber	9.55	10.04	8.87	8.32
Ether Extract	4.28	4.52	5.28	5.45
Ash	8.42	7.12	7.90	6.42
Nitrogen free extract	62.38	63.14	62.44	64.18
Acid detergent fiber	35.80	36.03	35.14	36.05
Neutral detergent fiber	12.94	13.02	13.22	12.65
Lignin	2.82	2.77	2.73	2.75
Cellulose	10.12	10.25	10.49	9.90
Hemicellulose	22.86	23.01	21.92	23.40
Metabolic energy (Mica Gul/kg)	12.3	12.5	12.6	12.8

Metabolic energy (Mg / kg of material as is) = 0.012 × crude protein + 0.031 x ether extract + 0.005 × raw fiber + 0.014 × nitrogen-free extract (Maff, 1975).

Table 6: Components of vitamins and minerals mix.

Vitamins	Concentration	Minerals	Concentration
Vitamin A	200 000 IU/kg	200 000 IU/kg	2000 mg/kg
Vitamin D3	100 000 IU/kg	100 000 IU/kg	2500 mg/kg
Vitamin E	515 mg/kg	515 mg/kg	1000 mg/kg
Vitamin B1	125 mg/kg	125 mg/kg	25 mg/kg
Vitamin B2	500 mg/kg	500 mg/kg	30 mg/kg
Vitamin B3	1000 mg/kg	1000 mg/kg	1200 mg/kg
Vitamin B6	35 mg/kg	35 mg/kg	1000 mg/kg
Vitamin B12	10 mg/kg	10 mg/kg	qsp mg/kg
		200 000 IU/kg	1500 mg/kg
		100 000 IU/kg	2000 mg/kg

Chemical analysis

The chemical analyzes of the feed samples were carried out, such as the untreated sunflower, the sunflower treated with blood, the sunflower treated with formaldehyde, dried whey powder treated with blood, the dried whey powder treated with formaldehyde, and the chemical analysis of the primary components of the experimental animals Table (1,4 &5) . These analyzes were carried out at the Central Laboratory of Graduate Studies, Nutrition Laboratory, Animal Production Department at the Faculty of Agricultural Engineering Sciences, University of Baghdad.

Study of some characteristics of the rumen

Samples of rumen fluid were extracted from 15 animals and 3 animals from each treatment during the eighth week of the experiment. The same animals were used in the digestion experiment where the rumen fluid was collected by a rubber tube inserted through the animal's mouth, reach the most appropriate point inside the animal's rumen to extract the rumen liquid through a 500 ml plastic syringe. The rumen liquid was withdrawn during three different times: the first time (zero), before the morning feeding. After this drawing, the concentrated feed was provided. The second and third withdrawn were taken after 3, 6 hours of morning feeding, respectively, and the rumen liquid was filtered with a piece of malmal to get rid of the solid particles immediately after the extraction of rumen fluid from the animal and converted to plastic bags to be saved, where 10 ml of rumen liquid was taken from each animal in each time of the three times (0, 3 and 6) hours of morning feeding. The pH of the rumen samples was measured and the bags containing the rumen liquid were divided into two parts. The first section was added 200 microliters (HCL) with the aim of stopping the microbial activity affecting the levels of both ammonia nitrogen and volatile fatty acids and then keeping these samples frozen until the analysis, while the samples of the other part of the folders containing the liquid rumen with a cooling temperature of 40C to study the total number of anaerobic bacteria also for the times when the rumen liquid was withdrawn (zero 3 and 6) hours of morning nutrition. Where the following measurements were made:

pH: After the rumen was collected directly, the pH was measured by the POCKET-SIZED PH METER (POCKET).

Nitrogen ammonia (NH₃-N): Nitrogen ammonia (NH₃-N - N) was estimated for rumen samples, according to A.O.A.C. (2005).

Total concentration of volatile fatty acids (TVFA): The total concentration of volatile fatty acids (TVFA) was estimated using the Kjeldahl distillation unit and the Warner method (1964).

Total count of anaerobic bacteria (BC): The total number of anaerobic bacteria was estimated by preparing the solutions and reagents, and the bacterial cultures were frightened. The media prepared according to what was stated in (Vucca, 1985; Atlas *et al.*, 1995; Vandepitte *et al.*, 2003) were used.

Dry matter DM: The dry matter of feed samples was estimated according to A.O.A.C. (2005).

Organic material (OM): Organic matter was calculated by subtracting the amount of ash from dry matter.

Crude protein CP: Crude protein was estimated using the Kjeldahl for fodder forms and according to A.O.A.C. (2005).

Crude fiber CF: Raw fiber was estimated for fodder models as indicated in A.O.A.C. (2005).

Ether Extract: The Ether extract for fodder samples was estimated according to A.O.A.C. (2005).

Carbohydrates dissolved in NFE: The dissolved carbohydrates were calculated according to the following equation: NFE = OM - (CP + CF + EE).

Neutral fiber extract: The NDF fiber extract was estimated according to Goering and Van Soest (1970).

Acid Fiber Extract: The acid fiber extract was estimated according to Goering and Van Soest (1970).

Acid fiber extract: The ADL extract was estimated according to Goering and Van Soest (1970).

Cellulose: Cellulose was calculated according to the following equation: Cellulose = ADF - ADL.

Hemicellulose: Hemicellulose was calculated according to the following equation: Hemicellulose = NDF- ADF.

Statistical analysis

The Statistical Analysis System (SAS) (2012) was used in data analysis to study Comparing the two experiences in the studied traits according to (Completely Randomized Design-CRD), The differences between the averages were compared with Test (T).

The mathematical model

$$Y_{ij} = \mu + E_i + e_{ij}$$

Y_{ij} = the value of the transaction j return to the transaction i.

μ = The general mean of the studied character.

E_i = It represents two experiences i.

e_{ij} = Random error distributed by a normal distribution with an average of 0 and a variance of σ^2e .

RESULTS AND DISCUSSION

There were no digestive disorders in the animals during and after the experiment period. All the animals were in a good health. The objective of the experiment was achieved by providing concentrated diets containing the ratio of sunflower treated with blood or formaldehyde, dried whey powder treated with blood or formaldehyde instead untreated sunflower with levels of (50 , 100%), while the green alfalfa was provided freely and the intake of concentrated feed, green alfalfa and total feed intake were calculated during the experiment period.

The rumen is an unstable fermentation vessel and according to the type of fermentation, so there are daily fluctuations in the Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N) and The total count of bacteria (BC), of the rumen (Emanuele & Putnam, 2006), while the level of protein degradation in the rumen is considered An important factor in influencing rumen variables that in turn affect the nature of digestion and microbial protein synthesis in ruminants, which are basic factors in animal need.

Effect type of protein (sunflower meal, dried whey powder) treatment with blood or formaldehyde on some Rumen Fermentations.

Table 7 showed that there were significant decrease (P < 0.05) in Rumen pH (pH) and insignificant effect in Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N) and The total count of anaerobic bacteria (BC), for diets dried whey powder treated with blood or formaldehyde compared diets sun flower meal treated with blood or formaldehyde. The reason for this may be Sun flower meal contains phenolic compounds that interact with formaldehyde before it interacts with the protein, which affects the level of protein protection from degradation compared to dried whey powder that does not contain phenolic compounds (Daiber & Taylor, 1982) This leads to an increase in the efficiency of protecting proteins from degradation in the rumen of diets dried whey powder treated with blood or formaldehyde compared diets sun flower meal treated with blood or formaldehyde (Parsons *et al.* 1991; Al- Shekhly, 1998). This leads to a decrease in the concentration of (NH₃-N) and an increase in the (TVFA) as in our current study of diets dried whey powder treated with blood or

formaldehyde (27,616) (11,733), respectively, compared to diets sun flower meal treated with blood or formaldehyde (28,390) (11,614), respectively, which leads to a lower pH of diets dried whey powder treated with blood or formaldehyde compared to diets sun flower

meal treated with blood or formaldehyde (Spears *et al.*, 1985; Moreira & Ribeiro 2000; - Al-Mallah2007; Shamoon *et al.*, 2009).

Fig 1. shows the effect of the type of protein treatment with (blood or formaldehyde) on some Rumen Fermentations of: Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N).




Fig. 1: Effect of the type of protein treatment with (blood or formaldehyde) on some Rumen Fermentations.

Table 7: Effect type of protein (sunflower meal, dried whey powder) treatment with blood or formaldehyde on some Rumen Fermentations

Studied traits	dried whey powder	standard error	sunflower meal	standard error	Effect significance
pH	6.502 ^b	0.070±	6.707 ^a	0.063±	*
TVFA (mmol /100 ml)	11.733	0.093±	11.614	0.094±	N. S
NH ₃ (mg/100 ml)	27.616	0.586±	28.390	0.632±	N. S
BC (cfu/ml)	210.72	10.145±	212.92	13.137±	N. S

Different characters within the same column indicate significant differences ($p < 0.05$); N.S Non-significant. Effect type of protein (sunflower meal, dried whey powder) treatment with blood in 100 % on some Rumen Fermentations.

Table 8 showed that there were no significant effect on the Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N) and The total count of bacteria (BC), for diets dried whey powder treated with blood in 100 % compared diets sun flower meal treated with blood in 100 % , these results are in

agreement with (Hadjipanayiotou, 1992 ‘saeed 2011 ‘Kahleefah, 2014).

Fig. 2 shows the effect of the type of protein treatment with blood in 100 % on some Rumen Fermentations of: Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N).




Fig. 2: Effect of the type of protein treatment with blood in 100 % on some Rumen Fermentations.

Table 8: Effect type of protein (sunflower meal, dried whey powder) treatment with blood in 100 % on some Rumen Fermentations.

Studied traits	dried whey powder	standard error	sunflower meal	standard error	Effect significance
pH	6.513	0.119 ±	6.660	0.110 ±	N. S
TVFA (mmol /100 ml)	11.998	0.172 ±	11.787	0.197 ±	N. S
NH ₃ (mg/100 ml)	25.920	1.133 ±	26.551	0.945 ±	N. S
BC (cfu/ml)	148.44	14.319 ±	201.67	31.923 ±	N. S

N.S Non-significant.

Effect type of protein (sunflower meal, dried whey powder) treatment with blood in 50 % on some Rumen Fermentations.

Table 9 showed that there were no significant effect on the Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N) and The total count of bacteria (BC), for diets dried whey powder treated with blood in 50 % compared diets sun flower meal treated with blood in 50 % , This may be due to the effect of treating dried whey powder and sunflower meal with blood or formaldehyde, as this led to significant

effects in the ratio of dry matter and the laboratory digestion factor of dry matter and organic matter, which led to the fact that the variation in their effects on rumen fermentations was not significant (Jasim 2019).

Fig. 3 shows the effect of the type of protein treatment with blood in 50 % on some Rumen Fermentations of: Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N).




Fig. 3: Effect of the type of protein treatment with blood in 50 % on some Rumen Fermentations.

Table 9: Effect type of protein (sunflower meal, dried whey powder) treatment with blood in 50 % on some Rumen Fermentations.

Studied traits	dried whey powder	standard error	sunflower meal	standard error	Effect significance
pH	6.677	0.117 ±	6.790	0.119 ±	N. S
TVFA (mmol /100 ml)	11.665	0.199 ±	11.560	0.185 ±	N. S
NH ₃ (mg/100 ml)	27.372	1.236 ±	28.218	1.620 ±	N. S
BC (cfu/ml)	214.56	17.248 ±	204.78	35.154 ±	N. S

N.S Non-significant.

Effect type of protein (sunflower meal, dried whey powder) treatment with formaldehyde in 100 % on some Rumen Fermentations.

Table 10 showed that there were no significant effect on the Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N) and The total count of bacteria (BC), for diets dried whey powder treated with formaldehyde in 100 % compared diets sun flower meal treated with formaldehyde in 100 % , these

results are in agreement with (Hadjipanayiotou, 1992 ; saeed; 2011 Kahleefah, 2014).

Fig. 4 shows the effect of the type of protein treatment with formaldehyde in 100 % on some Rumen Fermentations of: Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N).




Fig. 4: Effect of the type of protein treatment with formaldehyde in 100 % on some Rumen Fermentations.

Table 10: Effect type of protein (sunflower meal, dried whey powder) treatment with formaldehyde in 100 % on some Rumen Fermentations.

Studied traits	dried whey powder	standard error	sunflower meal	standard error	Effect significance
pH	6.400	0.169 ±	6.614	0.140 ±	N. S
TVFA (mmol /100 ml)	11.736	0.185 ±	11.583	0.195 ±	N. S
NH ₃ (mg/100 ml)	27.469	1.019 ±	28.227	1.228 ±	N. S
BC (cfu/ml)	242.89	16.947 ±	258.89	8.989 ±	N. S

N.S Non-significant.

Effect type of protein (sunflower meal, dried whey powder) treatment with formaldehyde in 50 % on some Rumen Fermentations.

Table 11 showed that there were no significant effect on the Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N) and The total count of bacteria (BC), for diets dried whey powder treated with formaldehyde in 50 % compared diets sunflower meal treated with formaldehyde in 50 % , This may be due to the effect of treating dried whey powder and sunflower meal with blood or formaldehyde, as this led to significant effects in the ratio of dry matter and the laboratory digestion factor of dry matter and organic matter, which led to the fact that the variation in their effects on rumen fermentations was not significant (Jasim 2019). The results of protecting protein from degradation in ruminant rumen may be conflicting due to the low level of protection in some cases and in other cases due to excessive protection as the protein becomes indigestible (Mir et al,1984). Many studies have shown

that a diet containing higher amounts of The non-dissolving proteins in the rumen or the amino acids protected from degradation in the rumen led to significant effects, while other studies showed little or no response, and the lack of response to the undegradable protein in the rumen is often due to one of the following reasons: 1- The undegradable proteins may be The rumen is bypassed the rumen at the expense of the rumen microbial protein synthesis. 2- The proteins that are undegradable in the rumen may be poorly digested after the rumen. 3- undegradable proteins in the rumen may be deficient in the amino acid content that limits production (Schingoethe, 1996).

Fig. 5 shows the effect of the type of protein treatment with formaldehyde in 50 % on some Rumen Fermentations of: Rumen pH (pH), Total volatile fatty acids (TVFA), Ammonia nitrogen in the rumen (NH₃-N).




Fig. 5: Effect of the type of protein treatment with formaldehyde in 50 % on some Rumen Fermentations.

Table 11: Effect type of protein (sunflower meal, dried whey powder) treatment with formaldehyde in 50 % on some Rumen Fermentations.

Studied traits	dried whey powder	standard error	sunflower meal	standard error	Effect significance
pH	6.420	0.156 ±	6.766	0.146 ±	N. S
TVFA (mmol /100 ml)	11.531	0.184 ±	11.526	0.196 ±	N. S
NH ₃ (mg/100 ml)	29.706	1.116 ±	30.568	0.974 ±	N. S
BC (cfu/ml)	237.00	17.589 ±	186.33	17.551 ±	N. S

N.S Non-significant.

REFERENCES

1. Emanuele, S.M. and Putnam, D. 2006. Encapsulating nutrients to improve reproduction and nitrogen utilization in ruminants. Ruminant Nutrition Symposium, Best Western Gateway Grand, Gainesville Florida. Feb. 1-2.
2. Dhali, A., D.P. Mishra, R.K. Mehla and S.K. Sirohi 2006. Usefulness of milk urea concentration to monitor the herd reproductive performance in crossbred karanfries' cows. *Asian-Aust. J. Anim. Sci.*, 19:26-30.
3. Nisa, M., A. Javaid, M.A. Shahzad and M. Sarwar 2008. Influence of varying ruminally degradable to undegradable protein ratio on nutrient intake, milk yield, nitrogen balance, conception rate and days open in early lactating Nili-Ravi buffaloes (*Bubalus bubalis*). *Asian-Aust. J. of Anim. Sci.*, 1.
4. Fébel, H. & Fekete, S. 1996. Factors influencing microbial growth and the efficiency of microbial protein synthesis: a review. *Acta Vet. Hung.* 44:39.
5. Ørskov, E.R. 1992. Protein Nutrition in Ruminants. 2nd Edition. Academic Press. New York, EUA.
6. Bach, A., Calsamiglia, S. & Stern, M.D. 2005. Nitrogen metabolism in the rumen. *J. Dairy Sci.* 88: 1.
7. Nolan, J.V. & Dobos, R.C. 2005. Nitrogen Transactions in Ruminants. In: Quantitative Aspects of Ruminant Digestion and Metabolism. 2nd Edition. CAB International. Wallingford, UK. p. 137.
8. Wallace, R. J., R. Onodera, and M. A. Cotta, 1997. Metabolism of nitrogen-containing compounds. pp. 283-328. In: The rumen microbial ecosystem. P.N. Hobson and C. S. Stewart (Eds.). 2nd ed. Chapman & Hall. London.
9. Strobel, H.J. and J.B. Russell 1986. Effect of pH and energy spilling on bacterial protein synthesis by carbohydrate-limited cultures of mixed rumen bacteria. *J. Dairy Sci.*, 69: 2941.
10. Cardozo, P., S. Calsamiglia, and A. Ferret. 2000. Effect of pH on microbial fermentation and nutrient flow in a dual flow continuous culture system. *J. Dairy Sci.* 83(Suppl. 1):265.
11. Cardozo, P., S. Calsamiglia, and A. Ferret. 2002. Effects of pH on nutrient digestion and microbial fermentation in a dual flow continuous culture system fed a high concentrate diet. *J. Dairy Sci.* 85(Suppl. 1):182.
12. Vucca, L. 1985. Laboratory manual of histochemistry Ravan press. New York. PP:328.
13. Atlas, R. M., L. C. Parks and A.E. Brown 1995. Laboratory Manual of Experimental Microbiology. Mosby-Yearbook, Tnc., Missouri.
14. Vandepitte, J, Verhaegen, J. Engback, K., Rohnerp, P., Piot, P. & Heuck, C. C. 2003. Basic Laboratory Procedure in Clinical Bacteriology. World Health Organization Geneva, Singapore.
15. Daiber, K.H. and J.R.N. Taylor 1982. Effect of formaldehyde on protein extraction and quality of high and low. Tanins sorghum. *J. Agri. Food Chem.* 30:70-72.
16. Parsons, C. M.; M.K. Hashimoto; K.J. Wedekind; and D.H. D.H. Baker 1991. Soybean protein Solubility in potassium Hydroxide an in vivo protein quality. *J. Anim Sci.* 69: 2918 – 2924.
17. Al-Shekhly M M (1998) Improvement utilization efficiency of sunflower meal as a feed for ruminant animals. *Master Thesis*. College of Agriculture, University of Baghdad.
18. Spears, J.W., J.H. Clark. And E.E. Hatfield 1985. Nitrogen utilization and ruminal fermentation in steers fed soybean meal treated with formaldehyde. *J. Anim. Sci.*, 60: 1072-1080.
19. Moreira, O.M.S.C. and J.M.C.R. Ribeiro 2000. Utilization of different concentrates by growing lambs. Ruminal degradation. CIHEAM - Options Mediterraneennes. 131-136.
20. Al-Mallah, O.D.M. 2007. Effect of protein levels in formaldehyde treated rations on coefficient of digestion and performance in Awassi lambs. Ph.D. Thesis, University of Mosul.
21. Shamoan, M., N. Saleh and N.Y. Abbo 2009. Effects of different levels of protein treated with formaldehyde on nutrients digestibility and some rumen and blood parameters in Awassi sheep. *Iraqi J. Vet. Sci.*, 23, Supplement II, 169-173.
22. Hadjipanayiotou, M., 1992. Effect of protein source and formaldehyde treatment on lactation performance of Chios ewes and Damascus goats. *Small Ruminant Res.*, 8: 185-197.
23. Mir, Macleod, G.K., Buelianan – Smith, J.G., D.G. Grieve and W.I. Grovum. 1984. Methods for protection soybean canola proteins from degradation in the rumen. *Can. J. Anim. Sci.*, 64:853-865.
24. Stock, R., T. Klopfenstein, D. Brink, R. Britton and D. Harmon. 1986. Whey as a source of rumen degradable protein. II. For Growing Ruminants production. *J. Anim. Sci.* 63:1574.
25. Hassan S A, Al- Ani A N, Al-Jassim R A M and Abdullah N S (1990) Effects of roughage to concentrate rations and rumen undegrable protein supplementation on growth of lambs. *Small Ruminant Research* 3, 317-324.
26. Kahleefah M M (2014) Effect of feeding Undegradable protein and bakery byproduct on Awassi lambs Performance. *Master Thesis*. College of Agriculture – University of Baghdad.
27. Cooper R J, Milton C T, Klopfenstein T J and Jordan D J (2002) Effect of corn processing on degradable intake protein requirement of finishing cattle. *J. Anim. Sci.* 80, 242-247.
28. Schingoethe, D.J. 1996. Balancing the amino acid needs of the dairy cow. *Anim. Feed Sci. Tech.*, 60: 153-160.
29. Dagher V J, Raz M A and Umayjan M (1980) Studies on the utilization of full fat sunflower seed in broiler ration. *Poult. Sci.* 59, 2273-2278.
30. USDA-FAS (2017) *Oilseeds: World Markets and Trade*. Available at: <https://apps.fas.usda.gov/psdonline/circulars/oilseeds.pdf>. Accessed Aug. 24, 2017.
31. Lusus E W C (1982) sunflower meals and food protein in sunflower Handbook National Sunflower Assn. Bismark, N.D. pp 26-36.
32. El-Shewy, A. 2016. Whey as a feed ingredient for lactating cattle. *Sci. Int.* 4: 80-85.
33. M. Ben Salem and M. Fraj 2007. The effects of feeding liquid acid whey in the diet of lactating dairy cows on milk production and composition. *Journal of Cell and Animal Biology* Vol. 1 (1), pp. 007- 010, August 2007.
34. Matsumoto M, Kobayashi T, Itabashi H and Ørskov E R (1995) Investigation of protection of soybean meal and amino acid from rumen degradation with whole blood treatment. *Anim. Feed Sci. Techn.* 56, 37-43.

35. MAFF. 1975. Ministry of Agric., Fisheries and Food Dept., of Agric. And Fisheries for Scotland Energy allowances and Feed systems for ruminants, Technical Bulletin,33. First published.
36. A O A C (2005) *Association of official analytical chemists, official methods of analysis*. 14th end. Washington, D.C., U.S.A.
37. Goering H K and Van Soest P J (1970) *Forage fiber analysis*, U.S. Department of Agriculture, Handbook. No. 379, 1-20.
38. SAS (2012) *Statistical Analysis System, User's Guide*. Statistical. Version 9.1st ed. SAS. Inst. Inc. Cary. N.C. USA.
39. Warner, A.S. 1964. Production of volatile fatty acid in the rumen methods of measurements. *Nutrition Abstract Review*, 34 pp. 339 – 343.
40. Saeed A A (2011) Effect of level and degradability of dietary protein fed with or without baker's yeast (*Saccharomyces cerevisia*) on Turkish awassi lambs' performance. *Ph.D* litter university of Baghdad, College of Agriculture.
41. Jasim, Ibrahim Sattar (2019) Effect of Substitution protected whey or sunflower meal with blood or Formaldehyde on Awassi lamb's performance. *PhD thesis*, College of Agriculture Engineering Science, University of Baghdad.