

Fast Food Consumption and Increased Caloric Intake Leading to Obesity a Survey among Pakistani Teenagers

Bilal Younas^{1*}, Waqas Ahmad Khalid², Mansoor Ul Hassan³

¹Rural Health Center, Lahore, Pakistan

²Rural Health Center, Raiwind, Lahore, Pakistan

³BHU 157 RB, Tehsil Chak Jhumra, Faisalabad, Pakistan

Article History:

Submitted: 12.04.2021

Accepted: 19.05.2021

Published: 26.05.2021

ABSTRACT

Background: The aim of the study was to determine obesity rate in Pakistani teenagers associated with fast food consumption.

Methods: This cross sectional study was conducted at RHC Raiwind, Lahore during January 2019 to July 2019. Information was collected from teenagers of different age groups ranging from 13-19 years old. A self-structured questionnaire was designed containing four sections to gather information from participants. BMI and % age of different variables was calculated.

Results: Total no of 1000 teenagers including male 536(53.6%) and females 464(46.4%) were approached out of which 334 (34.4%) were overweight having BMI=25-29.9 and 544(54.4%) were obese having BMI>30. Mostly of them were 17-year-old 425(42.5%).

Obesity rate is higher both in males and females but mostly females were more prone. Conclusion: It is concluded from our study that consumption of fast food on daily basis and increased caloric intake will leads towards obesity among teenagers now a days. Obesity rate is greater in females (refer to stage 4. of table. 3) as compared to males however both males and females were prone towards obesity due to increased consumption (thrice a day) of fast food. Other variables like residential area of participants and type of fast food eaten were also accountable in this regard.

Keywords: Obesity, Dietary habits, Glycemic load

***Correspondence:**

Bilal Younas, Medical Officer at RHC, Lahore, Pakistan, E-mail: bilalyounas37@yahoo.com

INTRODUCTION

The prevalence of overweight and obesity has also increased substantially in the nutritional transition countries, and the health burden of obesity-related complications is growing. The introduction of fast-food chains and Westernized dietary habits providing meals with fast-food characteristics seems to be a marker of the increasing prevalence of obesity. The mechanisms involved are probably that the supply of foods is characterized by large portion sizes with a high energy density, and sugar-rich soft drinks (Astrup *A et al.*, 2008). Fast food can be defined as convenience food purchased in self-service or carry-out eating venues without wait service (Pereira MA *et al.*, 2005).

In addition to the documented increase in fast food expenditures, many aspects of fast food make it suspect to the associated increases in overweight and obesity. Specifically, fast food tend to be energy dense, poor in micronutrients, low in fiber, high in glycemic load and excessive in portion size, causing many to exceed daily energy requirements (Pereira MA *et al.*, 2005) (Isganaitis E *et al.*, 2005).

Several dietary factors inherent to fast food may cause excessive weight gain such as massive portion size, high energy density, palatability (appealing to primordial taste preferences for fats, sugar, and salt), high content of saturated and trans fat, high glycemic load, and low content of fiber (French SA *et al.*, 2001).

The aims of this study were to determine whether fast food adversely affects diet quality in ways that might plausibly increase risk for obesity among teenagers.

An investigation was directed in Iran in which the national pervasiveness of overweight and heftiness, just as some related way of life practices, without precedent for Iran were assessed. Overall, 50.4% (n545 113) of the members were male and 64.6% (n557 866) were from the urban regions. The na-

tional appraisals of overweight, stoutness and bleak heftiness were 28.6%, 10.8% and 3.4%, separately. Weight list (BMI) \$25 kgm²² in men, ladies, urban inhabitants and provincial occupants were found in 37%, 48%, 46.7% and 35.5%, individually. Stomach corpulence was available in 43.4% of ladies, 9.7% of men, 28.5% of the urban inhabitants and 23% of the provincial occupants. Overweight just as summed up and stomach heftiness were increasingly common in the 45-64 years age gathering. In spite of the fact that there was no noteworthy distinction in recurrence of utilization of the nutrition classes in subjects with various BMI classifications, different sorts of physical exercises demonstrated a consistent decay with expanding BMI (Kelishadi R, 2008).

METHODOLOGY

This cross sectional study was conducted at RHC Raiwind, Lahore during January 2019 to July 2019. In which data was collected from total no of 1000 teenagers ranging from age of 13-19 years (Paeratakul S, 2003).

Study design

Data was collected from 1000 teenagers residing in Lahore, the most populated city of Pakistan. Information was gathered from teenager of different age groups ranging from 13-19 years old. Participants residing in different area of Lahore Pakistan were invited to take part in our survey. We asked them a set of question written in our questionnaire, containing both open and closed ended questions (Mushtaq MU *et al.*, 2001).

Data collection tool

A self- made questionnaire was used to gathered information from participants, consisting of four sections (Bowman SA *et al.*, 2004) (Davis B *et al.*, 2009).

Section 1. It contains informed consent.

Section 2. Containing information regarding participant ID, gender, age, weight, height, residential area.

Section 3. Containing food intake questions e.g. total no of meal per day including breakfast, lunch, snacks, dinner and fast food intake per day either once, twice or thrice, weekly or monthly. Also included question about type of fast food (burger, pizza, shawarma, chicken wings, Bar B Q, chicken sigi, roast, pratha rolls, pastas, potato fries, soft drinks etc)

Section 4. BMI calculation.

According to WHO criteria classified in 4 stages.

Stage1. Underweight if BMI < 18.5

Stage2. Normal if BMI = 18.5-24.9

Stage3. Over weight (pre obese) if BMI = 25-29.9

Stage 4. Obese if BMI > 30

Analysis

Percentage of fast food intake among male and female participants and their BMI were calculated by using SPSS version 20.

RESULTS

A total no of 1000 respondents (teenagers) were approached. Total no of male and females in our study were 536(53.6%) and 464(46.4%) respectively. Different age groups participants ranging from 13 to 19 years old were take part in study. Majority of them were 17 years old (42.5%). Other percentage of respondent's age groups given in Table 1.

Table 1: Demographic characteristics of respondents N=1000

Demographic variables	Frequency %
Gender	
Male	536 (53.6 %)
Female	464 (46.4 %)
Age (y)	
13-14	86 (8.6 %)
15-16	245 (24.5 %)
17	425 (42.5 %)
18-19	244 (24.4 %)

BMI calculations were given in Table 2. According to that 37 (3.7%), 85 (8.5%), 334 (33.4%), 544 (54.4%) fall in stage 1,2,3 and 4 respectively, which shows increased fast food consumption will lead obesity among teenagers. Also Table 3 gives detailed figures between BMI stages and other variables.

Table 2: BMI stages, frequency and percentages of respondents N=1000

Stages	Frequency	Percentage
Underweight if BMI < 18.5	37	3.7 %
Normal if BMI = 18.5-24.9	85	8.5 %
Over weight (pre obese) if BMI = 25-29.9	334	33.4 %
Obese if BMI > 30	544	54.4 %

Table 3: BMI stages among other variables

BMI Stages	Stage1	Stage2	Stage3	Stage4	Total
Gender					
Male	0	83	247	206	536
Female	37	2	87	338	464
Age (years)					
13-14	7	17	25	37	86
15-16	10	21	71	143	245
17	12	36	145	232	425
18-19	8	11	93	132	244

Fast food consumption					
Once a day	0	17	63	73	153
Twice a day	0	0	90	133	223
Thrice a day	0	0	125	271	396
weekly	5	23	42	54	124
Monthly	32	45	14	13	104
Residential area					
Near restaurants	13	33	281	460	787
Far restaurants	24	52	53	84	213
Type of fast food					
pizza	3	13	65	113	194
Burger	5	10	54	105	174
shawarma	2	15	45	97	159
Chicken wings	3	10	25	34	72
Paratha rolls	7	9	20	31	67
Pastas	2	5	17	12	36
Potato fries	1	3	29	37	70
Soft drinks	10	2	32	57	101
Bar B Q	2	2	14	33	51
Chicken sigi	1	9	13	12	35
Roast	1	7	20	13	41

DISCUSSION

In this survey we found alarming situation regarding the obesity rate among teenagers. Fast food consumption becoming a fashion and easy way to eat quickly prepared food items among teenagers. Other factors like palatability and easily availability of these items also involved. In our survey we found both male and female participants were prone to obesity but mostly females (338) falls in stage 4 of BMI (refer to table 3). In accordance to the age group mostly of our participants of 17 year old were obese.

Following three issues should need to be addressed.

Firstly consumption of fast food on daily basis also in mostly cases three times a day among teenagers causes increased obesity rate.

Secondly residential area play vital role in eating habits of fast food, participants living near by the fast food restaurants would like to prefer fast food on home made dishes and in our findings having more obesity rate as compared to those which are not living near restaurants.

Thirdly type of fast (pizza, burger, shawarma, and soft drinks) consumption is higher and leads obesity as compared to other items.

CONCLUSION

This study was conducted to find out the consumption of fast food and obesity rate among teenagers. Total no of 1000 participants take part in our study. Obesity rate is greater in females (stage 4 table 3) as compared to males. No male was fall in stage 1, but majority (247) were in stage 3 of BMI. And we concluded from our study that fast food consumption in teenagers is becoming a trend and its harmful effects will lead towards obesity. Mostly of participants were 17-year-old obese (42.5%). Fast food consumption on daily bases mostly thrice a day and residential area (near restaurants) and type of fast food (pizza, burger, shawarma, soft drinks) were majorly involved in obesity rate.

REFERENCES

1. Astrup A. Nutrition transition and its relationship to the development of obesity and related chronic diseases. *Obes Rev* 2008; 9: 48-52.

2. Pereira MA. Fast-food habits, weight gain, and insulin resistance (the CARDIA study): 15-year prospective analysis. *Lancet*. 2005; 365(9453): 36-42.
3. Isganaitis E, Lustig RH. Fast food, central nervous system insulin resistance, and obesity. *Arteriosclerosis, thrombosis, and vascular biology*. 2005; 25(12): 2451-2462.
4. French SA. Fast food restaurant use among adolescents: associations with nutrient intake, food choices and behavioral and psychosocial variables. *Int J obes*. 2001; 25(12): 1823.
5. Kelishadi R. Obesity and associated lifestyle behaviours in Iran: findings from the first national non-communicable disease risk factor surveillance survey. *Public health nutrition*, 2008. 11(3): 246-251.
6. Paeratakul S. Fast-food consumption among US adults and children: dietary and nutrient intake profile. *J Am dietetic Asso*. 2003; 103(10): 1332-1338.
7. Mushtaq MU. Dietary behaviors, physical activity and sedentary lifestyle associated with overweight and obesity, and their socio-demographic correlates, among Pakistani primary school children. *Int J Behav Nutri Phys Act*. 2011; 8(1): 130.
8. Bowman SA. Effects of fast-food consumption on energy intake and diet quality among children in a national household survey. *Pediatrics*. 2004; 113(1): 112-118.
9. Davis B, Carpenter C. Proximity of fast-food restaurants to schools and adolescent obesity. *Am J Pub Heal*. 2009; 99(3): 505-510.